



## International Journal of Current Research and Academic Review

ISSN: 2347-3215 Volume 2 Number 6 (June-2014) pp. 188-196

[www.ijcrar.com](http://www.ijcrar.com)



### Self-immolation Oriented Behaviors in young Militaries – A Review article

Zahra Amirmohamadi<sup>1</sup>, Abdollah Soltaninejad<sup>2\*</sup>, Ali Reza Nikmorad<sup>3</sup>,  
Hediye Sadat Mirsharafoddini<sup>4</sup>, and Mahrokh Bagheri Moghadam<sup>5</sup>

<sup>1</sup>Master of Entrepreneurship, Jiroft University of Medical Sciences, Jiroft, Iran

<sup>2</sup>Behavioral Sciences Research Center, Baqiyatallah University of Medical Sciences, Tehran, IR Iran

<sup>3</sup>Tehran Delaram Psychiatric Center, Tehran, Iran

<sup>4</sup>Department of Economic, Firoozkooh Branch, Islamic Azad University, Firoozkooh, Iran

<sup>5</sup>Department of Laboratory Science, North Tehran Branch, Islamic Azad University, Tehran, Iran

\*Corresponding author

#### KEYWORDS

Self-immolation,  
Self-immolation  
oriented  
behavior,  
self-mutilation,  
Police and  
military  
soldiers

#### A B S T R A C T

Self-immolation oriented behavior is one of the most important problems in mental health field. self-mutilation, suicide thinking, Self-immolation attempt and complete Self-immolation are behaviors that investigated in psychology as oriented behavior. Military and police are Self-immolation subject due to long term encounter to stress, adaptation problems, access to weapons and distinct position. The aim of this review article is investigation of different aspect of Self-immolation oriented behavior in police and military. This article is a review study that prepared with research in library source using reliable scientific documents and article. Some parts of this article are results from author PhD thesis in Self-immolation oriented behavior. Self-immolation is not a sickness, but it is the sickness outcome and the consequence of cooperation between biological, psychological, social and organizational traumas. Self-immolation and its related behaviors have destructive effect on survivors, victim's family, comrades and commanders. Self-immolation is a preventable event and with a suitable council, it can be control. The aim of Self-immolation prevention program is to minimize Self-immolation oriented behavior as low as possible. Screening and identification of personnel with Self-immolation risk, attend to soldier mental health, stress management in unit and encouragement of help-seeking

### Introduction

Self-immolation oriented behavior is a world spread problem and there are evidence for its global enhancement(1). Also there aren't indication for reduction of its related behaviors such as self-

mutilation that make competent a person for Self-immolation. Self-immolation oriented behavior has wide range and includes any think and action that shows a person is going to hurt himself. Self-

immolation thinking, Self-immolation attempt and complete Self-immolation are some example of this behaviors. Self-immolation thinking and previous Self-immolation attempt are two major factors in Self-immolation possibility. Self-immolation oriented behavior has complex pathology and includes wide range of familiar, biological, psychological, social, cultural and spiritual factors. Military forces due to access to weapons, distinct situation, young age and high level of stress are subject of Self-immolation risk. Self-immolation oriented behavior in any form and level has destructive effect on people, survivors and managers of mental health. Studies showed that family relative of Self-immolation military have depression symptoms, Self-immolation thinking and Posttraumatic Stress Disorder (2-3). Among Self-immolation oriented behavior, Self-immolation is a outstanding mental hurt in militaries that led to great damage for military forces .Identification of Self-immolation oriented behavior's reasons, incidence and its preventer factors has important role in design and implementation of precautionary programs.

### **Self-immolation oriented behavior in psychological literature**

Although in psychology and psycomedicine Self-immolation oriented behavior debate is focused on Self-immolation and self-mutilation, but suicidology science, especially in recent decade, emphasizes on Self-immolation oriented behavior as an actions that have common trauma and ultimately led to similar outcome such as death. When we speak about Self-immolation oriented behavior, we noticed to wide range of actions which include actions and thinks that can be conclude that a person is going to harm or Self-immolation himself .there

is other view about Self-immolation oriented behavior definition. This vision divided SOBs into 2 categories the first one is death tendentious and other is life tendentious SOBs. In is death tendentious SOBs that include Self-immolation attempt and complete Self-immolation, a person either died due to Self-immolation or his /her predisposition for Self-immolation is evaluated high(4). In life tendentious SOBs the person is not going to Self-immolation himself, but repetition of this actions directs person to the Self-immolation. self-mutilation and repetitive self-mutilation are two sample of life tendentious SOBs. There aren't any purpose for Self-immolation in self-mutilation. Also in repetitive self-mutilation that shows a person is encountered to very stressful situation, there e is not intend for Self-immolation too, however repetition of this action prone to person for Self-immolation.(5) Researchers that studied in SOB field are believed that SOBs are in opposite side of non SOBs. This action are distinct from non non SOBs due to existence of Self-immolation intent (6,7).

### **Epidemiology of SOBs in Iran and world**

Self-immolation is an important problem in public health in world spread. About one million people were died due to Self-immolation in 2000. Self-immolation is one of the 10 leading cause of death in public population and is the second or third cause of death in 15-34 years olds(8). The global rate of Self-immolation is 16 in 100000, in previous decade it is estimated that disease burden due to Self-immolation was 1.8%. Recently disease burden due to Self-immolation is growing and this rate possibly reaches to 2.4 in 2020. In past half century Self-immolation rate has grown about 60% (9). Based on WHO estimation about 1530000 people will die in 2020 due

to Self-immolation and 10 to 20 fold will have Self-immolation attempt(10).

Review of SOBs in Iran has heterogeneous results due to cultural and economical and social variation, and also different methodology. The review of 19 sequential Self-immolation in Iran By Ghoraihi et al showed that the age range of operators are 19-29 and the average of Self-immolation attempt is 39 and 61% in men and women(11)based on this study results the rate of Self-immolation in Iran is 9.4 in 100000 that is lower than global rate. The results of this study showed that the rate of men Self-immolation incidence in Hamedan, Ilam and Lorestan is higher than other regions. Also women Self-immolation in Ilam, Kermanshah and Lorestan is higher than other province. The common method for Self-immolation in Iranian is hanging and self-burning in Men and women respectively. This study declare wide spectrum of variables as Self-immolation risk factor including low income, low teaching, no logical believed, mental disorders especially behavior disorder, addiction, history of previous Self-immolation, married situation and gender. Although there are not clinical evaluation in 70% of SOBs studies, but there is a significant relation between mental disorder and Self-immolation attempt so that 45% of Self-immolation attempts have mental disorder(11).

### **Epidemiology of SOBs in militaries**

Self-immolation in military personnel is an important mental health issue due to the access to weapons and familiarity with its use, high stress and positions distinct from others. Although military forces have a proper mental health and physical fitness and are mentally assessed during the service period, they are at the risk of Self-

immolation or self-mutilation because of risk factors for Self-immolation (young age, access to weapons, stress, certain positions and aggressive people in military forces). Some experts believe that Self-immolation in military forces should be less than the general population due to screening and access to mental health services for soldiers (12), but there is another viewpoint against this one and claims that high aggression and access to guns increase the risk of Self-immolation in military forces. It seems that some evidences support the second view (13). Self-immolation is considered as the second cause of death in America and one of the leading causes of death in Russian army. Self-immolation in the United States military is constantly rising. This statistics has approximately increased from 9.6 per hundred thousand people in 2004 to 21.9 per hundred thousand people in 2009 and has approximately reached 22.9 per hundred thousand people between 2009 and 2010. Mental health status, problems related to substance abuse, problems associated with physical health such as cancer or chronic pain and problems related to financial, legal and communication issues have been identified as risk factors for Self-immolation among urban populations and military personnel (14). Studies have shown that over the past decade, the rate of Self-immolation among United States military has increased by 71% and reached from 9.1 per hundred thousand people in 2001 to 15.6 per hundred thousand people in 2010 (15). Studies by mental anatomical methods to identify risk factors in Yugoslav Army during 1998-2007 have shown that being single, heritable psychiatric disorders, ineffectiveness of psychiatric treatments, heavy physical exercise, not giving transfer to the other posts, low motivation to serve in the military profession, incompleteness

military education and the indebtedness of households are regarded as the risk factor for Self-immolation (16). In a field study among American navies and marine forces it was found that the ratio of Self-immolation leading to death to failed Self-immolation attempt is one-seventh among navies and one-fifth among marine forces (17-18).

The rate of Self-immolation attempts for women was two to three times more than men. In 95 % of psychological disorder cases, it was diagnosed that the main diagnoses were personality disorder (53%), followed by substance abuse (36 %). Study in United States Army Air Force during 1990-1994 has shown that twenty-three percent of all deaths among Air Force active personnel have been Self-immolation. The results of a study on the Polish armed forces in the period 1989-1998 have shown that the number of completed Self-immolations has been 437 cases in these ten years that among these, 163 cases have occurred in officers and 273 cases in soldiers (19).

There is no exact number of Self-immolations in Iranian armed forces, but unofficial statistics indicate the incidence of 9 per hundred thousand people. Studies of Anisi and colleagues on the prevalence of suicidal thoughts in ground force soldiers have shown that the incidence of suicidal thoughts among soldiers is 5.8 percent (4). The results of this study have shown that higher prevalence of Self-immolation thoughts is related to being single, drug abuse, and previous history of Self-immolation, Self-immolation in family, primary and guidance education, lack of mental health, serving in border areas and serving in service and security jobs (4). The results of Noori about the risk factors for Self-immolation in the soldiers

of Sepah Ground Forces have shown that the most important risk factors for soldiers' Self-immolation include previous Self-immolation attempt, depression, addiction, conflicts with work environment, family problems, aberrant humiliation and punishment, service in line units, the presence of personality traits such as introversion, neurosis and psychotic disorder (20). The results of Soltaninezhad and colleagues on 1659 soldiers in one of the Armed Forces in country's six provinces in 2012 have shown that the incidence rate of Self-immolation oriented behaviors including suicidal thoughts and the history of Self-immolation attempts is 15.9% in this population (4). Studies of Roohani and colleagues on 321 employees in NEZAJA garrison in Tehran have shown that 15.3 % of studied sample had suicidal thoughts and approximately 1.9 % of them had a history of Self-immolation attempts. Based on the results of this review, 5.9 % of participants have reported a history of psychiatric disease. This review has shown that the prevalence of suicidal thoughts has been 19.8 % in draftees and 84.2% in soldiers with psychiatric disorder (21). The study of Matinsadr and colleagues on 327 soldiers has shown that 15.1 % of studied sample have suicidal thoughts and 8.4% have the history of one to seven Self-immolation attempts (22). The study of Soltaninezhad and colleagues on 1659 soldiers of one of the Armed Forces in country's 6 provinces has shown that the prevalence of suicidal thoughts in the sample is 10.9 % and the prevalence of Self-immolation attempts 5.7% and total incidence rates of Self-immolation-oriented behaviors in this group is 15.9% (23).

### **Risk factors for Self-immolation**

Numerous studies around the world have identified several risk factors for Self-

immolation-oriented behaviors. Although in these studies, that each one has been done with specific objectives, have been used different methods, there is a high agreement on identifying risk factors for Self-immolation-oriented behaviors (24, 25, and 26).

According to the International Health, Hemmati and colleagues have categorized and presented specific risk factors associated with a high risk of Self-immolation in three categories.

These factors are:

1. Demographic factors: male gender, age between 25-44 years old, living in the countryside, being members of minority groups, refugees, homeless people and sexual identity conflict

2. Groups at risk of Self-immolation: history of previous Self-immolation attempts, previous self-mutilation, history of mental illness, especially depression, schizophrenia and other psychotic disorders, personality disorders, history of sexual abuse, drug or alcohol abuse, or both, diseases or severe physical disabilities, imprisonment or detention

3. Recent risk factors: having psychological characteristics such as helplessness, hopelessness, confusion, shame, feelings of guilt, shyness, psychotic thoughts, interpersonal conflicts, being excluded from a particular person, the recent loss of someone or something, feeling bad luck or misfortune and recent unemployment, alcohol consumption, chronic and painful disease, financial difficulties, being fired and recent unemployment, to commit a crime, disintegration of the family, custody of a family member, lack of social support networks, reluctance to seek help and lack of access to mental health services.

Self-immolation Society of America has identified and introduces five categories of risk factors for Self-immolation. The

factors are stable and non-adjustable risk factors, adjustable and predisposing risk factors, auxiliary risk factors, acute risk factors and starter risk factors (28).

### **Stable and non-adjustable risk factors of Self-immolation**

-Demographic factors including being white, male gender, ageing, divorce, separation, early widowhood

-Previous suicidal thoughts, history of repeated Self-immolation attempts, history of self-mutilation, history of Self-immolation in family

-History of injury or physical or sexual abuse, history of psychiatric hospitalization, history of repeated relocation, history of violent behaviors and history of impulsive behaviors.

### **Adjustable and Predisposing Risk Factors**

-Psychiatric disorders axis 1, especially anxiety disorders, schizophrenia, drug or alcohol abuse, eating disorders, body dysmorphic disorders, conduct disorder

-Disorders axis 2 including personality disorders, especially personality disorders cluster B

-Disorders axis 3, especially medical disorders, especially if causes malfunction or be associated with chronic pain, minor brain injuries

-Being accompanied of disorders axis 1, 2, and 3.

### **Auxiliary Risk Factors**

-Easy access to weapons, unemployment, stress in the field of occupation, marriage, school, interpersonal relationships

### **Acute Risk Factors**

-Recent divorce or separation, especially if it is accompanied with the feel of anger and being victim.

-Suicidal thoughts, Self-immolation plan, threat to Self-immolation or being ready for Self-immolation

-Recent self-mutilation, recent Self-immolation attempt

-Indulgence in substance abuse

-Psychological pain (severe distress following exclusion, loss or failure)

-Anger, wrath, aggressive behavior, seeking for revenge

-Lack of enjoyment, anxiety, restlessness, pain, insomnia, frequent or persistent nightmares,

-Imperative illusions with the subject of insistence on Self-immolation

-Severe emotional moods such as depression, self-blame, unbearable loneliness, excessive mood changes

Starter and Detector Risk Factors, conditional on personal vulnerability to Self-immolation

-Any event that causes shame, frustration and feelings of guilt a

-To stand accused, conflict with the law, custody, business failures, being excluded from an important person

-To witness the death of a friend or relative who had died of Self-immolation or watch the same events in media

### **Self-immolation Prevention**

Self-immolation and its related behaviors can be prevented. The aim in Self-immolation prevention is to minimize the Self-immolation oriented behaviors. There are several main strategies in Self-immolation prevention that play a pivotal role in Self-immolation prevention plans. These strategies include: promoting mental health level and prevention of mental health problems, treatment of patients with mental disorders, especially those with mental disorders related to Self-immolation oriented behaviors, restriction of access to weapons and lethal tools and also

strengthening skills to deal with mental health problems, especially skills in seeking help from others in dealing with mental health problems and also religious coping skills (29-35). Self-immolation prevention programs in military personnel are also affected by national strategies for Self-immolation prevention in countries. Self-immolation prevention programs in the United States Air Force are mental health awareness and to encourage help-seeking behaviors, to attract the participation of commanders and staff in the field of Self-immolation prevention, to train mental health professionals, to establish Self-immolation-oriented behaviors research center, to launch intervention teams in traumatic crisis and Self-immolation and to identify Self-immolation risk factors. The results of evaluating the Self-immolation prevention programs have shown that about a third of those whom the program is applied, have shown decrease in domestic violence, homicide and Self-immolation risk factors (36). Self-immolation prevention program in the United States Army is based on four overall strategies. The first strategy for Self-immolation prevention is based on identifying those at risk for Self-immolation. Screening soldiers to identify those who are at risk of Self-immolation is the first step in implementing Self-immolation prevention programs. The second strategy is to attract the participation of commanders and managers, especially commanders and leaders who are at the first line of communication with soldiers. According to this strategy, commanders and managers should be well trained about the risk factors and protective factors of Self-immolation and be able to identify those at risk and refer them. Mental health administrators should consider identifying Self-immolation starters and symptoms of people involved in crisis as an important issue in training

commanders and managers. The third strategy is to promote and disseminate the culture of asking for help. All commanders should encourage soldiers and their subordinates to seek help without fear of its consequences. Feeling guilt, shame, embarrassment and fear of disciplinary or retaliation actions are serious obstacles for soldiers on the way to seek help. The fourth strategy is to increase the mental strength and resilience through coping skills training.

### **Discussion and Conclusion**

Self-immolation-oriented behaviors are affected by individual, family and social, physiological, psychological, cultural, economic and spiritual factors (30, 34 and 390). In addition to the risk factors known in the etiology of Self-immolation in the general population, soldiers youth, poor coping skills, economic and family problems, characteristics of the military environment, aggression and access to guns, the discount rate for officers, mental disorders, dissatisfaction with the military, poor quality relationship with commander are among the factors playing a role in suicidal behaviors in the military forces (28, 38 and 41). Self-immolation is preventive, provided that Self-immolation prevention plans to be based upon the pathology of trauma and developed by taking into account the cultural and social factors and unit's capabilities and following the multi-dimensional approach.

### **References**

- 1.Columba Mclaughlin (2007). Self-immolation Related Behavior. John Wily& Sons, Ltd.
2. Helena Korosec Jagodic, Mark Agius & Peter Pregelj. INTER REGIONAL VARIATIONS IN SELF-IMMOLATION RATES. *Psychiatria Danubina*. 2012 ; 24(1) : 82–85.
- 3.Jong-Min Woo, Olaoluwa Okusaga and Teodor T. Postolache. Seasonality of Suicidal Behavior. *Int J Environ Res Public Health* 2012 ; 9 (3) : 531-547.
- 4.Anisi et al . Prevalence and factors associated with suicidal ideation among soldiers. *Journal of Military Medicine* , 2008 ; 8 (2) , 113-118.
- 5.Edward A. Selby, Michael D. Anestis, Theodore W. Bender, Jessica D. Ribeiro, Matthew K. Overcoming the fear of lethal injury: Evaluating suicidal behavior in the military through the lens of the Interpersonal–Psychological Theory of Self-immolation. *Clinical Psychology Review*. 2010 ; 30(7) : 298–307.
- 6.Mary Jane Alexander,Gary Haugland, Peter Ashenden, Ed Knight. Coping With Thoughts of Self-immolation:Techniques Used by Consumersof Mental Health Services. *Ps psychiatryonline org*. 2009 ; 60(9) : 121-125.
- 7.Joseph Logan. Nancy A. Skopp,. Debra Karch, , Mark A. Reger, and Gregory A. Gahm, Characteristics of Self-immolations Among US Army Active Duty Personnel in 17 US States From 2005 to 2007. *American Journal of Public Health* . 2012 ; 102(1) : 56-60.
- 8.Yanhong Gong, Ling Zhang, , Zhiqing Wang, Yuan Liang. Pathway Analysis of Risk Factors for Severe Suicidal Ideation;A Survey in Rural China. *J Public Health* 2011;102(6):472-75.
- 9.World Health Organization,(2008). Available from [www.WHO.com](http://www.WHO.com)
- 10.Jose Manoel Bertoolote,and Alexander Fleischmann. A global Perspective in the epidemidemiology of Self-immolation. *Suicidologi* 2002 arg ,7.nr.2.

11. Qureshi A et al . Systematic review of studies of Self-immolation attempt in Iran. *Iranian Journal of Psychiatry and Clinical Psychology* . 2008 ; 14 (2) : 115-121.
12. Richard J. Pinder, Amy C. Iversen, Nav Kapur, Simon Wessely and Nicola T. Fear. Self-harm and attempted Self-immolation among UK Armed Forces personnel: Results of a cross-sectional survey. Downloaded from [isp.sagepub.com](http://isp.sagepub.com) at Kings College London - ISS on June 22, 2011.
13. Charlotta Sunnqvist<sup>1</sup>, Lil Träskman-Bendz<sup>2</sup>, Åsa Westrin. Coping strategies used by Self-immolation attempters and comparison groups. *Open Journal of Psychiatry*. 2013; 3(2): 256-263.
14. American Foundation Fore Self-immolation Prevention(2011).available [www.afsp.org](http://www.afsp.org).
15. Judy Regan and et al. Self-immolation in the Military. *TENNESSEE MEDICINE / JULY 2005*.
16. Dedić G, Panić M. Self-immolation risk factors in the professional military personnel in the Army of Serbia] *Vojnomedicinska akademija, Klinika za psihijatriju, Odeljenje za mentalno zdravlje I vojnu psihologiju, Beograd, Srbija. Vojnosanit Pregl*. 2010;67(4):303-12.
17. Drew M. Velting. Suicidal ideation and the Five-factor model of personality. *Personality and Individual Differences*. 1999 ; 27(5) : 943-952.
18. Department of Health and Human Services (HHS). Substance Abuse and Mental Health Services Administration(2008), [www.samhsa.gov](http://www.samhsa.gov).
19. Fathi Ashtiani A, Salimy SH. Assessment of psychopathology and suicidal causes from 1989 to 1998 soldiers. *Journal of Military Medicine* , 2000 ; 3(4) : 249-254.
20. Nori R , et al. Important risk factor for Self-immolation among soldiers, a military force. *Journal of Military Medicine* . 2012 ; 14(2) : 99-103.
21. Rohani, M. et al . Prevalence of suicidal thinking in military barracks Army Tehran in 2004 . *Iranian Army Journal of Medical Sciences* ; 4(3) : 909-914.
22. Matini Sadr et al. Factors affecting group cohesion and their relationship to aggressive and suicidal thoughts draftees. *Journal of Military Medicine*. 2009 ; 11(4) : 237-242.
23. Soltani-Nejad, A., et al. Examine the relationship between religious orientation behavior Khvdksygrayanh soldiers. *Journal of Islamic lifestyle*. 2012 ; 1(2)32-37.
24. Arin N. Alberts . A Self-immolation Prevention and Intervention Program for Incarcerated Youth In the State of Illinois(2009).
25. Palmer, Stephan . Self-immolation Strategies and Intervention for Reduction and Prevention. published by Routledge. USA (2008).
26. Kimberly A. Van Orden; Tracy K. Witte,; Kelly C. Cukrowicz; Scott Braithwaite; Edward A. Selby. The Interpersonal Theory of Self-immolation. *Psychol Rev*. 2010 ; 117(2): 575–600.
27. Hemmati, et al . Build a scale to detect individuals at risk for the city of Mahabad and Bukan committed Self-immolation in 1386. *Quarterly Order and Safety* . 2009 ; 2 (1) : 29-58.
28. Jeffrey Hyman, Robert Ireland, , Lucinda Frost, and Linda Cottrell. Self-immolation Incidence and Risk Factors in an Active Duty US Military Population. *American Journal of Public Health*. 2012 ; 102(1) : 29-34.



29. LaRicka R, Wingate LR, Bobadilla L, Burns AB, Cukrowicz KC, Hernandez A, Ketterman RL. Suicidality in African American men: the roles of southern residence, religiosity, and social support. *Self-immolation Life Threat Behav.* 2005;35(6):615-29.
30. Cherise L. Wells . A Comparison of Suicidal Ideation between Active Duty Military and Military Dependents, University of the Rockies, Colorado School of Professional Psycholog(2008).
31. Rothmann, S et al. Psychological strengths, coping and Self-immolation ideation in the south african police services in the north west province. *Sa. Journal of Industrial Psychology.* 2002 ; 28(3) : 39- 49.
32. Teresa Kirchner, Laia Ferrer, Maria Forns, Daniela Zanini. Self-harm behavior and suicidal ideation among high school students. Gender differences and relationship with coping strategies. *Actas Esp Psiquiatr.* 2011;39(4):226-35.
33. Andrej Marusic, MRCPsych, MD, PhD, and Renee D. Goodwin . Suicidal and Deliberate Self-Harm Ideation Among Patients with Physical Illness: The Role of Coping Styles. *Self-immolation and Life-Threatening Behavior.* 2006 ; 36(3) : 56-60.
34. Department of Health and Human Services (HHS). Substance Abuse and Mental Health Services Administration(2008), [www.samhsa.gov](http://www.samhsa.gov) v.
35. Abdollah Soltaninejad et al . Personality Factors Underlying Suicidal Behavior Among Military Youth. *Iran Red Crescent Med J.* 2014 April; 16(4): e12686.
36. National Strategy for Self-immolation Prevention. Goals and Objectives For Action. A report of the U.S. Surgeon General and of the National Action Alliance for Self-immolation Preventio2012.
37. Army Health Promotion Risk Reduction & Self-immolation Prevention(2010).
38. Harold Braswell , Howard I. Kushner. Self-immolation, social integration, and masculinity in the U.S. military. *Social Science & Medicine.* 2012 ; 74(6) : 536-539.
39. Yari G and Alan A. Self-immolation and Suicidal Behavior. *Public Health Reviews.* 2012 ;34(2) : 33-38.
40. Soltani-Nejad, A., et al. Examine the relationship between religious orientation and Self-immolation attempt soldiers. *Journal of Military Psychology .* 2012 ; 2(1) : 23-29.
41. Soltani-Nejad, A., et al. Structural model of the relationship between borderline personality disorder, coping style, emotion, impulsivity and suicidal thoughts soldiers. *Medical Journal of Police .* 2012 ; 1(3) : 176-182.