




International Journal of Current Research and Academic Review

ISSN: 2347-3215 Volume 3 Number 11 (November-2015) pp. 190-198

www.ijcrar.com



Study on Genetic diversity and relationship of silkworm varieties using RAPD molecular marker

B.K.Chikkaswamy and Rabin Chandra Paramanik*

Sigma Bioscience Research Centre, 2nd stage, Indiranagar, Bangalore-560038, India

**Corresponding author*

KEYWORDS

Genetic diversity,
Silkworm varieties,
Molecular marker,
RAPD

A B S T R A C T

The genomic DNA of eleven silkworm were amplified with OPA-01 primer revealed 102 RAPD bands out of which 80% is polymorphism and 20% is monomorphic silkworm is polygenic. Some of the DNA fragments were strain specific and some could differentiate the multivoltine from the bivoltine strains or vice versa. Silkworm genetic resources that are being maintained in Hosur, Tamil Nadu, India, are yet to be adequately topped to develop elite varieties that are subjected to different agro-eco-climatic condition of the country like India. Molecular markers are known to provide unambiguous estimated of genetic variability of silkworm populations. Since they are independent confounding effects of environment, the genetic similarity among the seventeen silkworm varieties/strains and their genetic diversity and relationships were discussed.

Introduction

Mulberry silkworm (*Bomby mori* L.), the most important silk producing insect, exhibits wide diversity in morpho-biochemical and biometric characters. In general, silkworm strains from temperate countries like China, Japan and Korea complete two generations per year (bivoltines) by adopting egg diapause. On the other hand, those from tropical countries like India, Bangladesh and tropical belts of China complete five to six generations, per year without undergoing diapause, hence they are called polyvoltine silkworms. The polyvoltines are comparatively smaller in

size but can with stand adverse climatic conditions such as higher humidity, temperature and to a certain extent exhibit tolerance to diseases. However, from the productivity point of view, bivoltines are preferred'. Since the bivoltines are prone to abiotic and biotic stresses prevailing in the made to develop silkworm hybrids with productivity traits of bivoltines and tolerance of polyvoltines. In order to achieve this goal, knowledge on the genetic make up of silkworm strains is necessary. The present investigation is an attempt in this direction

where bivoltines are compared with polyvoltines along with a few mutants derived from bivoltines but with intermediate voltinism pattern (Table 1). Using RAPD markers, researchers have characterized silkworm strains. However, in the present study the silkworm mutants, which are believed to have originated from bivoltines, but closer to polyvoltines in yield attributes, are also included to understand the genetic relatedness with either of the genotypes.

The ISSR primers are core sequences of SSRs and reported to be widely spread over genome, provide more realistic genetic information in closely related genotypes like silkworm. In addition to RAPD and ISSR markers, RFLP-STS markers are also used, thus including both dominant and co-dominant markers for drawing precise conclusions on their genetic relatedness.

The biology and genetics of silkworm, *Bombyx mori*, is the most advanced of any lepidopteran species. Its rich repertoire of genetic resources and potential applications in sericulture and as a model for other Lepidoptera led to the initiation of genomics research. During the past decade much effort has been made in the areas of marker development, and molecular maps have been constructed in standard strains with the use of RFLPs, RAPDs, ISSRs, STSs, and microsatellites. The potential applications of molecular markers and linkage maps include stock identification, Marker Assisted Selection (MAS), identification of Quantitative Trait Loci (QTL), and, ultimately, positional cloning of visible mutations and QTL. To these ends, BAC libraries have been constructed and are being used to make large-scale physical maps, with markers based on ESTs as framework anchors. Altogether this work provides a foundation for identification of

gene function, gene and chromosome evolution, and comparative genomics.

Genetic Markers

In the past decade, several key advances in molecular biology have greatly advanced the impact of molecular genetics in biology. Most important have been: (i) development of the polymerase chain reaction (PCR), which amplifies specific stretches of DNA to usable concentrations, (ii) the application of evolutionarily conserved sequences as PCR primers, (iii) the advent of hyper variable microsatellite loci as genetic markers and (iv) the advent of routine DNA sequencing in biology laboratories. These innovations, coupled with the recent explosion of powerful analysis and relatively user-friendly computer programs, have accelerated the pace at which molecular genetic data can be used for various programmes. These developments led to the detection, analysis and exploitation of naturally occurring DNA sequence polymorphisms in eukaryotes and laid a foundation for contemporary genomics of several organisms. Polymorphic genetic markers have found potential applications in animal and plant improvement programmes as a means for varietal and parentage identification, construction of molecular linkage maps and evaluation of polymorphic genetic loci affecting quantitative economic traits.

A wide array of DNA marker techniques is available for genetic studies. All DNA markers reflect differences in DNA sequences. Typically, in a diploid organism, each individual can have one or two different states (alleles) per character (locus). The choice of a particular genetic marker often depends upon the purpose of the study and is usually a trade-off between practicality and precision of genetic

markers. One manifestation of this is the dichotomy between multilocus DNA techniques which include Random Amplified Polymorphic DNAs (RAPDs), Amplified Fragment Length Polymorphisms (AFLPs) and Inter-Simple Sequence Repeat PCR markers (ISSR-PCR) and single locus techniques which include single copy nuclear DNA regions and microsatellites. Multilocus methods result in simultaneous visualization of many dominant markers and are technically convenient, but have some marked weaknesses and limitations, including dominant inheritance - the DNA fragment can be scored only as present or absent, and in many cases substantial proportion of variation they detect can be non-heritable or not even derived from the target organism. By contrast, single locus markers are quite informative because they provide alleles whose zygosity status can be assayed easily.

Considering the genetic informativeness offered by microsatellite loci, which are co-dominant, a microsatellite discovery programme in the silkworm genome was initiated. Microsatellites, also known as simple sequence repeats (SSRs), are short stretches of nucleotides in which a motif of one to six bases is tandemly repeated, have emerged as ideal markers in eukaryotic genetics. They are ubiquitous in prokaryotic and eukaryotic genomes and are randomly distributed, both in protein coding and non-coding regions. The sequences in microsatellite motifs can differ in repeat number among individuals, providing a ready source of polymorphism. With the advent of the PCR, this property of microsatellite DNA was converted into highly versatile genetic markers. PCR products of different lengths can be amplified using primers flanking the variable microsatellite region. Due to the advantages offered by microsatellites they

have been used to construct linkage maps. In addition to their clear utility for practical applications, information about the distribution and variability of microsatellite sequences in the genome of a given species can help elucidate its genetic history from the standpoint of evolution and artificial selection.

The perusal of the earlier literature indicates that most of the RAPD and ISSR molecular markers were carried out on different varieties silkworm. It is also noted that earlier researchers have not investigated the molecular markers studies in some of the silkworm varieties. The investigation of RAPD is necessary to be used as a tool to investigate the genetic diversity and relationships of silkworm varieties. It is appropriate to evaluate their use in breeding purpose. Although sporadic reports on molecular markers in silkworm varieties (are mostly known from the earlier researchers) they have not examined the RAPD and ISSR in detail in other silkworm varieties regarding it is, there fore pertinent to investigate the above said parameters

In the present investigation, ten silkworm varieties from Hosur germplasm Tamil nadu, was selected to carryout RAPD molecular markers to study genetic diversity and relationships silkworm varieties

Materials and Methods

The present investigation was carriedout in the Department of Om –Bioscience research center, 2nd stage, Indiranagar, Bangalore, and Horticulture, Biotechnology Centre, DNA Fingerprinting Laboratory, Govt of Karnataka, Hulimavu, Banerghata Road, Bangalore.Karnataka.India, during the year January-2006 to December-2007.The materials used and methods followed in this study are presented here.

Silkworm Strains

The ten different silkworm strains viz Sanish E2(M), Sanish E2(M), Sanish-8, Sanish-17, Sanish-18(M), Sanish-18(P), Sheikhi, Sanih-30, Sheikhi-II and Tetra Hybrid (P) comprising of bivoltine and polyvoltine accessions silkworm stocks having divergent geographical origin were used in the study. These silkworms were maintained at Central Sericultural Germplasm Resources Centre, Hosur, Tamil Nadu, India (77.51°E, 12.45°N, 942m AMSL) [www.silkgermplasm.com], following the recommended rearing practices.

DNA Isolation

Genomic DNA was extracted from the silkworm moth using phenolchloroform method briefly; Silk moths were frozen first with liquid nitrogen and homogenized with cooled mortar and pestle. The powdered content was transferred to an Oakridge tube and 5 mL of the extraction buffer (100 mM Tris-HCl, 8.0,50 mM EDTA and 1% SDS) and 25 uL of proteinase K (100 ug/mL) were added before incubating it at 37°C for 2 h with occasional swirling. DNA was extracted twice with phenol-chloroform-isoamyl alcohol (24:24:1) and once with chloroform.

The supernatant DNA was precipitated in ethanol in the presence of 3 M sodium acetate (pH 5.2), re-suspended in TE buffer (10 mM Tris-HCl, 1 mM EDTA, pH 8.0). The RNA contamination was removed by incubating with RNase A (100 ug/mL) at 37°C for 1 h. DNA was re-extracted with phenol-chloroform and precipitated with ethanol. The genomic DNA was quantified on 0.8% agarose gel and diluted to uniform concentrations (10 ng/ul) for PCR amplification.

PCR Amplification of Genomic DNA

For RAPD, the amplification of genomic DNA was carried out in a PTC-200 Thermal-cycler (MJ Research Inc., Watertown, Massachusetts, USA) using 12 random decamer primers will be obtained from Operon Technologies, USA (OPW series) following the standard procedure Twenty-one ISSR (Inter simple sequence repeat) primers will be procured from the University of British Columbia, Vancouver, Canada (UBC set No. 9), will be tested for their efficacy in amplification of Silkworm DNA employed in this study. PCR amplification of the DNA was carried in the Thermal-Cycler, PTC 200¹⁰.

DNA Electrophoresis

Amplified DNA fragments were separated out on 1.2% agarose gel stained with ethidium bromide. Running buffer containing Tris-buffer, Acetic acid and EDTA (pH 8.0) was used for electrophoresis and for preparing gels. Wells were loaded with 25µl reaction volume and 5 µl of loading buffer (Sucrose, Bromophenol blue and Xylene cyanol) together. Electrophoresis was conducted at 45 volts for 3 hours and the gel photographed under UV light using gel dock system (Herolab).

Statistical Analysis

Binary coding was used to score gel and each band of primer was scored of 6 species and 12 primers with 100 to 1000 base pairs Marker level pair wise squared Euclidean species was calculated and utilizing these distances, species were clustered following Wards method. The segregation of species was also assessed through principle component analysis, the Statistica version 5.0 a computer application was used to

generate dendrogram using squared Euclidean distance and Wards method.

Results and Discussion

The yield and quality of DNA extracted by following the procedure described earlier was 10-25 ng per μ l for silkworm moth. The DNA obtained was amplifiable and of high quality. Spectrophotometer reading of 1.7-1.8 (260nm/280nm) confirmed the quality of DNA. DNA isolated from silkworm moth using 20 ml extraction buffer yielded good quality, high molecular weight DNA (above 50 kb). The quality of the DNA was also confirmed by gel electrophoresis.

In this study, 100 Operon random ten-base long, single stranded arbitrary primers (OPW -02, OPW-O4, OPW-05 OPW-06 OPW-011 OPW-12 OPW-13 OPW16 OPW17 OPW18 OPW19 and OPW-20 Table-1) were screened using the silkworm varieties, which on an average gave nine bands. The selected 12 arbitrary RAPD primers were used for the screening of the silkworm varieties, in which 20% monomorphic and 80% polymorphic and were considered for the precise calculation of genetic diversity. Twelve RAPD primers revealed 102 markers of 82 which polymorphic. All the silkworm varieties grouped in to four groups and indicating RAPD bands lie in between the 1.0 kb to 2.0kb

The basic protocol described elsewhere for PCR was optimized for high quality amplification and intense repeatable banding patterns. However, a reduction in the amplification of fainter bands was noticed with large changes in template DNA concentrations, while too much DNA produced a smear effect, which emphasized the importance of quantification of the DNA for clear amplification.

A representative of the PCR amplification product of the 10 silkworm varieties is as shown in fig.1 which yielded sufficient polymorphisms to distinguish between the silkworm varieties by Ward's method of analysis with Squared Euclidean distance gave dendrogram. Based on the number of bands all the 10 silkworm varieties were grouped in three clusters (**Fig. 2**). The dendrogram revealed a maximum similarity between the silkworm varieties. Every single silkworm accessions /hybrid could be identified using these selected twelve primers.

Analysis of diversity based on PCR fragments amounted to saturating the genome. The analysis of 10 silkworm varieties revealed that diversity is moderate to high and has shown even differences also. The maximum polymorphism between silkworm varieties could reveal significant correlation.

The data regarding RAPD molecular marker on selected 10 silkworm varieties showed considerable variability. The larval duration was in the different ranges and final instar duration varied from from 128-181 h. The larval, cocoon and shell weight were showing considerable variation between silkworm varieties. The squared Euclidean distances was calculated using the data on 10 silkworm varieties with 12 Primers (OPW -02, OPW-O4, OPW-05 OPW-06 OPW-011 OPW-12 OPW-13 OPW16 OPW17 OPW18 OPW19 and OPW-20 Table-1) Highest distance was observed between JD and Sanish E2 and Sanish E2 and SH-6. The lowest distance was recorded between Sheikhi-1 and Sheikhi-2. The distance within silkworm varieties was least (2.84 units) and highest (4.6 units) and intermediate distance between (3.82 units). However, this was in line with the results obtained in Wards genetic diversity

calculations. The dendrogram (Fig.2) showed four distinct clusters namely Cluster1 Sanish E2a, SanishE2b, Sanish8 and Cluster 2 included Sanish17, Sanish18, Sanish30, Cluster3 Sheikhi-i Sheikhi-ii tetra hybrid (p) kyorieshimpaku (p) kyorieshimpaku (m) and shinki reyaku (m0) and cluster4 included NB4D2 Sf-19 Sh-6 and Jd-6 From the pattern of clustering, it was pertinent that RAPD technique was efficient in segregating silkworm varieties into different four clusters. More significantly, the clustering had been largely successful in retaining the relationship between Silkworm varieties as proposed by B.K.Chikkaswamy In the light of current study at varietal level, it can be seen in the clustering pattern that the series were clearly distinguished in silkworm varieties. The association of the varieties observed in the present study was similar to the pattern observed by Awasthi et.al, (2008) and Nagaraj and Nagaraj (1995).and Chatterjee and Pradeep (2003).

Differential Polymorphism of DNA

The result presented in the present study demonstrated the utility of using RAPD markers to characterize genetic diversity among 10 silkworm varieties. Differential polymorphism was noted in 10 silkworm varieties showing variation in percentage of polymorphic bands from 20% to 80% using 12 Primers (OPW -02, OPW-O4, OPW-05 OPW-06 OPW-011 OPW-12 OPW-13 OPW16 OPW17 OPW18 OPW19 and OPW-20 Table-1) (. The observed high proportion of polymorphic loci reveals profound variation among the silkworm varieties. Significant genetic variations by RAPD have also been reported in other species level Awasthi et.al, (2008), Nagaraj and Nagaraj (1995).and Chatterjee and Pradeep (2003). Basavaraja, et al (2001), Benchamin, et al (1988), Bhandari, et al (1997), Bhargava et al (1995), Buteler and

LaBaonte (1999), Chandel and Joshi (1983), Chatterjee, and Datta (1992), Chatterjee et al (1990), Chatterjee et al (1993), Chattopadhyay et al (1995), Chatterjee (1993), Das et al (1987), Das et al (1994), Data and Ashwath (2000), Data (1984), Data and Basavaraja, (1994) and Data and Pershad, (1988) Wide genetic distances determined by Euclidean Wards genetic distance reveals relatively high genetic variation among 10 silkworm varieties. The considerable polymorphism detected in the present day also illustrated genetic diversity among silkworm varieties of the same origin as reported among other varieties of Silkworm (by Srivastava(2005). The observed differences among 10 silkworm varieties could be ascribed to the fluctuating micro and macro climatic conditions of habitat. Ward's analysis of RAPD data also reveals that all silkworm varieties belonging to the state of Karnataka are genetically closer. The greater sensitivity of RAPD obtained in the result to species diversity may be derived from rapid evolution of non-coding, repetitive DNA sequence detected by RAPD. This hypothesis has been corroborated from (Plomion et al. 1995). Akagi et al (1996), Akagi, et al. (1997), Anand, . and Murthy, (1968) , Anand, and Rawat, (1984), Arunachalam, and Bandyopadhyay, (1984), Arunachalam, Bandyopadhyay, Nigam, and Gibbons, (1984), Ashoka, and Govindan, (1990a), Ashoka, and Govindan, (1990b), On comparison of 10 silkworm varieties in, it can be observed that though they differ largely in their, phenotypic expression, but the genetic difference as observed from DNA fingerprints was found to be low. The genetic variation as detected by RAPD analysis opens up the avenue for the proper identification and selection of the species/varieties that could be used for varietal identification and planning for future breeding programme.

Table.1 Name, Origin and Quantitative Traits of 10 Silkworm Accessions Studied

SL.No.	TLD (h)	VLD (h)	LWT (g)	CWT (g)	SWT (g)	SR (%)
1	585	151	37.26	1.700	0.298	17
2	620	177	37.437	1.686	0.339	20.189
3	609	170	35.498	1.662	0.320	19.288
4	632	182	39.098	1.655	0.336	20.384
5	628	182	38.914	1.714	0.356	20.857
6	626	180	40.089	1.752	0.331	19.014
7	558	147	24.585	1.163	0.173	14.973
8	551	143	27.068	1.257	0.191	15.383
9	574	169	26.294	1.368	0.191	14.104
10	550	143	21.071	1.039	0.143	13.920

Table.2 Primers and Percentage of Polymorphism Realized with the RAPD Primers

SL.No.	Primer (Operon)	Sequence	No. of total bands	No. of polymorphic bands	% polymorphism
1	OPW-02	ACCCCGCCAA	8	7	90.00
2	OPW-04	CAGAAGCGGA	8	7	90.00
3	OPW-05	GGCGGATAAG	8	7	90.00
4	OPW-06	AGGCCCGATG	8	5	70.48
5	OPW-11	CTGATGCGTG	8	7	90.00
6	OPW-12	TGGGCAGAAG	10	7	70.00
7	OPW-13	CACAGCGACA	9	7	80.00
8	OPW-16	CAGCCTACCA	8	7	90.00
9	OPW-17	GTCCTGGGTT	8	7	90.91
10	OPW-18	TTCAGGGCAC	9	6	80.89
11	OPW-19	CAAAGCGCTC	9	8	80.00
12	OPW-20	TGTGGCAGCA	9	7	90.91
Total/Avg			102	82	93.60

Figure.1 RAPD Profiles of Genomic DNA of 10 Silkworm Accessions Generated with Random Primers OPW-02





Figure.2 Dendrogram Based on RAPD Marker Using Euclidian Distance Computer Version 0.5



RAPD analysis revealed a high degree of genetic diversity among the species/varieties, examined in the study, which can contribute to the silkworm crop involvement. RAPD analysis can also be used for detecting gene flow between species. Furthermore, this technique is less restricting than other molecular technique RFLP (Restricted Fragment Length Polymorphism), as no hybridization and no use of radioisotopes is required, and it is therefore more convenient for use in research centers in developing countries. Thus our results demonstrate that DNA content and RAPD markers provide an effective tool for detection and evaluation of genetic variation existing among 10 silkworm varieties of the species *Bombyx mori*. Seventeen diverse strains of Silkworm (*Bombyx mori*) were analysed using the sequence repeat polymerase chain reaction for RAPD molecular marker.

References

- Akagi, H.; Yokozeki, A.; Nakamura, A. And Fujimura, T. (1996) A codominant DNA marker closely linked to the rice nuclear restore gene, Rf-1, identified with inter-SSR fingerprinting. *Genome*, 39: 1205-1209.
- Akagi, H.; Yokozeki, Y; Inagaki, A and Fugimura, T. (1997) highly polymorphic micro satellites of rice consist of AT repeats and a classification of closely related cultivars with these micro satellite loci. *Theor. Appl. Genet.*, 94: 61-67.
- Allard, R.W. (1960) Principles of plant breeding. John Wiley and Sons Inc., New York.
- Anand, I.J. and Murthy, B.R. (1968) Genetic divergence and hybrid performance in linseed. *Ind. J. Genet. P1. Breed.*, 28: 178-185.
- Anand, I.J. and Rawat, D.S. (1984) Genetic diversity, combining ability and heterosis in brown mustard. *Ind. J. Genet.*, 44: 226-234.
- Arunachalam, V. and Bandyopadhyay, A. (1984) Limits to genetic divergence for occurrence of heterosis-experimental evidence from crop plants. *Indian J. Genet.*, 44(3): 548-554.
- Arunachalam, V.; Bandyopadhyay, A.; Nigam, S.N. and Gibbons, R.W. (1984) Heterosis in relation to genetic divergence and specific combining ability in groundnut (*Arachis hypogaea* L). *Euphytica*, 33: 33-39.
- Ashoka, J. and Govindan, R. (1990a) Genetic estimates for quantitative traits in bivoltine silkworm, *Bombyx mori* L. *Mysore J. Agric. Sci.*, 24(3): 371-374.

- Ashoka, J. and Govindan, R. (1990b) Heterosis for pupal and related traits in single and double cross hybrids of bivoltine silkworm, *Bombyx mori* L. *Entomologia*, 15: 203-206.
- Basavaraja, H.K.; Kumar, N.S.; Reddy, N.M.; Kalpana, G.V.; Kumar, S.N.; Jayaswal, K.P.; Joge, P.G.; Palit, A.K. and Datta, R.K. (2001) New system of maintenance and multiplication of CSR breeds. *Hid. Silk*, 40(2): 5-10.
- Benchamin, K.V.; Jolly, M.S. and Benjamin, D.A.I. (1988) Studies on the reciprocal crosses of multivoltine and biovoltine breeds in silkworm *Bombyx mori* L. with special reference to the use of biovoltine hybrids as parent. *Hid. J. Seric.*, 27(1): 27-34.
- Bhandari, m.M.; Gupta, G.S. and Gupta, R. (1997) Genetic divergence in opium poppy (*Papaver somniferum* L.) *J. Genet.*, 57(1): 11-13.
- Bhargava, S.K.; Majunclar, M.K. and Datta R.K. (1995) Combining ability of seven silk technological characters in mulberry silkworm, *Bombyx mori* L. *Bull.Zool.*, 62: 359-362.
- Buteler, Jarret R.L. and LaBaonte, D.R. (1999) Sequence characterization of micro satellites in diploid and polyploidy *Ipomoea*. *Theor. Appl. Genet.*, 99: 123-132.
- Chandel, K.P.S. and joshi, B.S. (1983) Multivariate analysis in green seeded pea. *Ind. Agric. Sci.*, 53(4): 198-200.
- Chatterjee, S.N. (1993) silkworm breeding in India. *Sericologia*, 33(3): 427-447.
- Chatterjee, S.N. and Datta, R.K. (1992) Hierarchical clustering of 54 races and strains of the mulberry silkworm, *Bombyx mori* L.; Significance of biochemical parameters. *Theor. Genet.*, 5(4): 394-402.
- Chatterjee, S.N.; Nagaraj, C.S. and Giridhar, K. (1990) an approach to silkworm breeding. Workshop Biometrical genet, C.S.R.T. & 1, Mysore, 7-8 September, 11-16.
- Chatterjee, S.N.; Rao, P.r.M.; Jayaswal, K.P., Singh, R. and Datta, R.K. (1993) Genetic variability in mulberry silkworm, *Bombyx mori* L. Breeds with low silk yield. *Ind. J. Seric.*, 32: 69-86.
- Chattopadhyay, S.; Das, S.K., Roy, G.C.; Sen, S.K. and Sinha, S.S. (1995) Heterosis analysis on silk productivity of three way crosses in *Bombyx mori* L. *Sericologia*, 35(3): 549-551.
- Das, B.C.; Trag, A.R.; Shaheen, A.; Nabi, G. (1987) new hopes for sericulture in Kashmir. *Ind. Silk*, 25(9): 25.
- Das, S.K.; Patnaik, S.; Ghosh, B.; Singh, T.; Nair, B.P.; Sen, S.K. and Subba Rao, G. (1994) Heterosis analysis in some three way crosses of *Bombyx mori* L. *Sericologia*, 34(1): 51-61.
- Data R.K. and Ashwath, S.K. (2000) Strategies in genetics and molecular biology for strengthening silkworm breeding. *Ind. J. Seric.*, 39(1): 1-8.
- Data, R.K. (1984) Improvement of silkworm races (*Bombyx mori* L.) in India. *Sericologia*, 24(3): 393-415.
- Data, R.K. and Basavaraja, H.K. (1994) silkworm breeding in Japan, china and India a comparative analysis. In: *Silkworm breeding, Proc. Nat. Workshop, March 18-19*, Ed. By G.S. Reddy, Oxford and IBH Publ. Co., Pvt., Ltd., New Delhi, 18-38pp.
- Data, R.K. and Pershad, G.D. (1988) Combining abilities among multivoltine x biovoltine silkworm (*bombyx mori* L.) hybrids. *Sericologia*, 28(1): 21-29.