



*International Journal of Current Research  
and Academic Review*

ISSN: 2347-3215 Volume 3 Number 3 (March-2015) pp. 74-79

[www.ijcrar.com](http://www.ijcrar.com)



**External Genitalic structures of two species of genus *Euplexia* stephens  
(Noctuidae: Lepidoptera)**

Jagbir Singh<sup>1</sup> and Charan Kamal Sekhon<sup>2\*</sup>

<sup>1</sup>Department of Zoology and Environmental Sciences, Punjabi University, Patiala-147001, Punjab, India

<sup>2</sup>Department of Zoology, Sri Guru Granth Sahib World University (SGGSWU), Fatehgarh Sahib, Punjab – 140406, India

\*Corresponding author

**KEYWORDS**

Moths,  
Lepidoptera,  
semifascia,  
conducta,  
genitalia

**A B S T R A C T**

Genitalic attributes are very important for authentic identification of any species. In the present communication external male and female genitalic attributes of *Euplexia conducta* Walker and *Euplexia semifascia* (Walker) have been studied and illustrated in detail for the concrete diagnosis of a species.

**Introduction**

The genus *Euplexia* was proposed by Stephens on type species *lucipara* Linnaeus in 1829. This large genus comprises large number of species distributed in many parts of the World, but in present study, only two species of this genus has been studied and illustrated in detail.

Four specimens of *Euplexia conducta* Walker and six specimens of *Euplexia semifasica* (Walker) have been collected from different localities of Western Ghats of India. Male and female genitalia have been discussed and photographed for the first time. A key based on external male genitalia has also been constructed.

**Materials and Methods**

The adult representatives of two Noctuid moth species were collected from the florescent lights fitted at different places in Western Ghats of India. The collected moths were killed and preserved in air tight wooden boxes. For the preparation of slides of external male and female genitalia, abdomen of preserved specimens were detached and potashed in 10% solution (Robinson 1976), washed in 1% glacial acetic acid and dissected in 30% alcohol for taking out male and female genitalia. After proper dehydration in different grades of alcohol, the genitalic structures were cleared in clove oil and then mounted in Canada balsam on cavity slides. The photography of external male and female genitalic structures

was done with the help of image processing unit in the department of Zoology, Punjabi University, Patiala. The terminology given by Klots (1970) has been followed in the present studies for nomenclature purpose. The adult Noctuid moths along with dissected specimens were preserved in insect cabinets.

## Observations

### Genus *Euplexia* Stephens

Stephens, 1829, *Nom. Br. Insects.*, 1829: 41.

**Type species:** *Euplexia lucipara* Linnaeus.

**Distribution:** Nearctic and Palaearctic regions; throughout India, Sri Lanka and Myanmar; Fiji.

**Diagnosis:** Eye naked, with or without lashes; proboscis fully formed; palpi upturned, the second joint not reaching vertex of head and fringed with hair, the third prominent. Thorax with a slight tuft of outspreading hair behind the collar and paired tufts on metathorax. Abdomen with dorsal tufts on proximal segments; tibial without spines. Forewing with the cilia crenulate.

### Key to the studied species of genus *Euplexia* Stephens

1. Male genitalia with uncus long, needle like; tegumen as long as uncus; juxta candle like; vesica with cornuti present . . . . . ***conducta* Walker**
- Male genitalia with uncus short, thick, sharp at tip; tegumen longer than uncus; juxta tongue shaped; vesica membranous with cornuti absent . . . . . ***semifascia* (Walker)**

### *Euplexia conducta* Walker

Walker, 1829, *Cat.*, 10: 296.

**Male genitalia:** Uncus long, curved, needle like, tip pointed, weakly sclerotized; paired socii present; tegumen broad, weakly sclerotized, as long as uncus, both the arms broad, inverted u-shaped; vinculum sclerotized, smaller than tegumen, u-shaped; saccus reduced; valvae broad, foot like, symmetrical, densely setosed with macro setae; costa broad, concave; sacculus sclerotized, slightly excurved beyond middle; harpe well marked, sickle shaped; juxta candle like; transtilla membranous; aedeagus broad, tube like, sclerotized distally; vesica membranous, partially

scobinate bearing two sclerotized patches having bristled cornuti; ductus ejaculatorious entering medially.


**Female genitalia:** Corpus bursae rounded, membranous with scobination, narrow and sclerotized towards ductus bursae; signum absent; ductus bursae narrow, short, partially sclerotized tube; ostium bursae simple; posterior apophyses almost of equal length as anterior apophyses; papilla analis oval, broad, sclerotized with macro and micro setae.

**Wing Span:** 36mm.

**Old Distribution:** Throughout India, Sri Lanka & Burma; Andamans; Fiji.




*Euplexia conducta* Walker



A. Male genitalia, B. Aedeagus, C. Uncus with Tegumen (Lateral view),  
D. Valva (Left), E. Female genitalia, F. Corpus bursae (Enlarged),  
G. Papilla analis with Apophyses (Enlarged)



*Euplexia semifascia* (Walker)



A. Male genitalia, B. Aedeagus, C. Uncus with Tegumen (Lateral view),  
D. Valva (Left)

### Material examined

Himachal Pardesh: Nauni: 27.ix.06, 2♀♀,  
Sabathu: 7.viii.06, 1♂♂,  
Barog: 18.viii.08, 3♂♂♂.

### *Euplexia semifascia* (Walker)

Walker, 1856, *Cat. Lep. Het.*, **33**: 737.

**Male genitalia:** Uncus strong, curved, narrow at base, broad medially with pointed apex, sparsely setosed with setae; tegumen longer than uncus, both the arms highly broad, inverted u-shaped; vinculum sclerotized, less broad, v-shaped, as long as tegumen; saccus present; valvae well developed, symmetrical differentiated into parts; costal margin excurved, sparsely setosed with setae at apex; harpe and ampulla well marked; sacculus sclerotized, developed into strong dentate projections; juxta tongue shaped; transtilla present, sclerotized; aedeagus long tube like, sclerotized; vesica membranous, partially curved with sclerotized patches; ductus ejaculatorious entering subapically.

**Wing Span:** 38mm.

**Old Distribution:** N. W. Himalayas; Satara; Nilgiris.

### Material examined

Himachal Pardesh: Kharapathar .  
21.vi.06, 2♂♂♂, Koti: 8.vii.08, 1♂♂,  
Vazula: 19.vi.08, 2♂♂♂.

### References

- Berio E. Diagnosi di Eteroceri Africini. Mem Soc Ent Ital Geneva 1940; 19:125-128.
- Berio E. Nuove specie e generi di Noctuidae Africane e Asiatiche e note sinonimiche. Part II. Annali del Museo Civico di Storia Naturale Giacomo Doria 1973; 79:126-171.
- Bethune-Baker GT. New Noctuidae from British New Guinea. Novit zool 1906; 13:191- 87.
- Bryk F. Zur Kenntnis der Grossschmetterlinge von Korea. II. Macrofrenatae Ark Zool 949; 41(1):1-225.
- Clarke JFG. The Lepidoptera of Rapa Island- Smithson. Conli Zool 1971; 56:I-IV +1-282.
- Draudt M. Beitrage zur Kenntniss der Agrotiden-Fauna Chinas. Aus den Ausbeuten Dr Hone's. Mitteil Munchn Ent Ges 1950; 49:1-174.
- Felder C, Rogenhofer M. Reise der österreichischen Fregatte Novara um die Erde in den Jahren 1857, 1858, 1859 unter den Behilfen des Commodore B. von Wüllerstorff-Urbair. Zoologischer Theil. Band 2.Abtheilung 2. Lepidoptera. Rhopalocera Reise Fregatte Novara (Inhalts-Verz.) 1874; 1-9 (pl. 1-74).
- Hampson GF. Catalogue of Lepidoptera-Phalanae in the British Museum. Cat Lepid Phalaenae Br Mus 1908; 7:692.
- Hampson GF. Description of New Heterocera from India. Trans Ent Soc London 1895, 277-315.
- Hampson GF. Illustrations of typical specimens of Lepidoptera Heterocera in the collection of the British Museum. Part 8. The Lepidoptera of Heterocera of the Nilgiri district III. Typical Spec Lep Het Colln Br Mus 1891; 8:1-144, pl. 139-156.
- Hampson GF. Illustrations of typical specimens of Lepidoptera Heterocera in the collection of the British Museum. Part 9. The macrolepidoptera heterocera of Ceylon III. Typical Spec Lep Het Colln Br Mus 1893; 9:1- 182.

- Hampson GF. The Fauna of British India, including Ceylon and Burma, Moths 2, Taylor and Francis Ltd., London, 1894, 1-609.
- Hampson GF. The moths of South Africa (Part II). *Ann S Afr Mus* 1902, 255-446.
- Hreblay M, Ronkay L. Noctuidae from Nepal. Haruta, T. (ed.): Moths of Nepal. Pt 5. *Tinea* 15 (Suppl. 1). 1998, 117-310, pis 144-157.
- Kononenko VS. A revised catalogue of types of the Noctuidae (Lepidoptera) described by F. Bryk (1948) from the Korean peninsula. *Inse Kore* 1996; 13:1-26.
- Laporte B. Nouvelles especes de noctuelle trifides Africaines (Lepidopteres). *Bulletin Mensuel de la Societe Linneenne de Lyon* 1977; 46:297-303.
- Leech JH. New species of Deltoids and Pyrales from Corea, North China, and Japan. *Entomologist* 1889; 22:62-71, pl. 2-4.
- Moore F. Descriptions of new Indian Lepidopterous Insects from the collection of the late Mr. W.S. Atkinson. *Descr Indian lep Atkinson* 1882; (2):89-198, pl. 4-5.
- Poole RW. *Lepidopterorum catalogus* (N. Ser.), Fasc. 118 Noctuidae. E J Brill Leiden 1989, 1-1314.
- Rothschild W. On Lepidoptera from the islands of Ceram (Seran), Buru, Bali, and Misol. *Novit Zool* 1915; 22(1):105-144.
- Stephens JF. The Nomenclature of British Insects; being a compendious list of such species as are contained in the International Journal of Multidisciplinary Research and Development Systematic Catalogue of British Insects, and forming a guide to their classification, London: Baldwin and Cradock, 1829, 1-68.
- Strand EH. Sauter's Formosa-ausbeute: Noctuidae II Nebst nachträgen zu den familien Arctiidae, Lymantriidae, Notodontidae, Geometridae, Thyrididae, Pyralidae, Tortricidae, Gelechiidae un Oecophoridae. *Archiv Naturg* 1920; 84(12):102-197.
- Sugi S. Notes on some Japanese genera of the Noctuidae with descriptions of new species (Lepidoptera). *Tinea* Tokyo 1958; 4:179-199.
- Swinhoe C. A list of Lepidoptera of the Khasia Hills. *Trans Ent Soc Lond* 1895, 1-75.
- Turner AJ. New Australian Lepidoptera. *Trans R Soc S Aust* 1902; 26:175-201.
- Viette PEL. Description de nouvelles especes de Noctuelles quadrifides (Lepidoptera: Noctuidae). *Mem Inst Sci Madag Tananarive (E)* 1957; 7:17-139.
- Walker F. Characters of undescribed Lepidoptera Heterocera. EW Johnson London, 1869, 1-112.
- Warren W. New oriental Noctuidae in the Tring Museum. *Novit Zool* 1916; 23:210-227.