



doi: <https://doi.org/10.20546/ijcrar.2024.1204.005>

## Determination of the Major Plant Nutrient Contents in Parthenium (*Parthenium hysterophorus* L.) Compost at Ginir District of Bale Zone, Oromia Region, Southeast Ethiopia

**Tesfaye Ketema Defar\***, Mulugeta Eshetu Diriba, Girma Getachew Gemechu and Regassa Gosa Telila

*Sinana Agricultural Research Center, Soil Fertility Improvement, Soil and Water Conservation and Watershed Management Team, P. O. Box 208, Bale-Robe, Ethiopia*

\*Corresponding author

### Abstract

The experiment was conducted in the Ginir District of the Bale Zone in the Oromia Region in the southeast of Ethiopia. Thus, the objective of this study was to evaluate the major plant nutrients in compost made from parthenium combined with wheat residue and farmyard manure, as well as its overall quality and nutrient contents. To fulfill the designated purpose, The parthenium plants were gathered before flowering and chopped into smaller pieces. Based on this, it can be concluded that the preparation of the compost and the material sources play a significant role, especially since different materials and methods were used throughout the study. The parthenium compost was thus made separately for each of the following three categories or treatments: parthenium biomass plus farm yard manure, parthenium biomass plus crop residue, and parthenium biomass combination with both farm yard manure and crop residue. After being prepared and harvested, the parthenium compost was subjected to a final laboratory analysis. Using conventional laboratory techniques, the main chemical properties, including pH, EC, OC, TN, available P, CEC, exchangeable bases (Ca, Mg, K, and Na), and micronutrients (Fe, Mn, Cu, and Zn), were measured. The obtained results for the nutrient content characterizations were: pH; Ec; OM; TN; CEC; 7.1 to 7.27; 0.000058 to 0.000062 ds/m; 35.2 to 37.8%; 1.83% to 1.98%; and 34.8 to 53.2 cmol+/kg, respectively. Exchangeable bases exhibit a similar trend for the major essential plant nutrients. The results indicated that the compost had a high concentration of plant nutrients and varied significantly between the three parthenium compost preparation methods. Parthenium compost, therefore, offers multiple benefits, including high nutrient contents, weed control capabilities, and generally environmentally sound uses of organic fertilizers.

### Article Info

Received: 20 February 2024

Accepted: 28 March 2024

Available Online: 20 April 2024

### Keywords

Total N, nutrients, Parthenium, Parthenium compost, environmental sound.

### Introduction

The aggressive alien weed species Parthenium (*Parthenium hysterophorus* L., Asteraceae) is native to America (Kohli *et al.*, 2006), but it has since spread widely throughout Asia, Africa, and Australia (Evans,

1997). In the 1970s, parthenium weed was first unintentionally brought to Ethiopia. In 1988, parthenium was discovered for the first time in Ethiopia, at Dire-Dawa in the east, and later in the northeast, close to Desse (Seifu, 1990). Major hubs for the distribution of food aid, it was thought that parthenium weed seeds were

brought in from subtropical North America to contaminate grain food aid during the famine of the 1980s (Tamado and Milberg, 2000). Later, it quickly spreads throughout the entire nation, along roads and railroads in grazing areas and on arable land, having a significant impact on biodiversity, crop production, and animal husbandry (Tefera, 2002). The central rift valley of Ethiopia, as well as the nearby areas of the Afar Region, East Shoa, Arsi, and Bale in southern Ethiopia, are currently home to a large parthenium population.

According to Anbalagan and Manivannan (2012) and Jelin and Dhanarajan (2013), composting could be a helpful substitute for converting this species' biomass into a material that could be utilized as a soil conditioner. Utilizing sustainable manures in agriculture is a component of organic farming. While production was increased by using more chemical fertilizers, the soil's fertility was decreased because there wasn't enough organic matter in the soil. Using organic materials is advised to counter this. A promising method for recycling wastes and weeds is composting, which produces a product that enhances crop productivity and soil fertility without endangering the environment. It is simple to use, safe for the environment and helps with pollution issues (Yadav, 2015). Composting is a waste management technique that has been around for at least a century, and it is currently gaining attention from all over the world for its ability to reduce the amount of accumulated waste and use weeds (Yadav, 2015). In addition to competing with pasture and crop species, parthenium weed has been linked to human and animal health risks (Devi *et al.*, 2014). Crop growth and development can be inhibited, if not completely controlled, by parthenium. Farmers in the Bale zone refer to it as "Anamalee," which means "Only me" in Afaan Oromo, because of its aggressive coverage (Personal Communications). As per various authors (Wabuye *et al.*, 2014; Ayele *et al.*, 2014; Kumari *et al.*, 2014), parthenium is a species that is spreading and has a significant impact on biodiversity, agriculture, and natural ecosystem production. According to several studies (Ameta *et al.*, 2016; Fitsum *et al.*, 2017), parthenium compost possesses twice as much nitrogen, phosphorus, and potassium as farm yard fertilizer, making it useful for both weed eradication and organic fertilizer supplies.

Despite an abundance of locally accessible parthenium weed and a sufficient quantity of various essential macro and micro plant nutrients, farmers in the study area do not compost parthenium. Furthermore, very few, if any,

scientific studies have been done on the application of parthenium as a compostable material and its potential for better crop production instead of its eradication. To specifically characterize the quality and nutrient contents of compost made from parthenium combined with wheat residue and farmyard manure in terms of major plant nutrients, this study was conducted.

## **Materials and Methods**

As part of the Bale highlands in the Oromia Regional State in southeast Ethiopia, the study was carried out in the Ginnir District. At 07° 15' N latitude and 40° 66' E longitude, and 1972 meters above sea level, Ginnir is located 519 kilometers away from Addis Ababa (figure 1). There is 531 mm of seasonal rainfall and 13.4 and 25.5°C for mean annual minimum and maximum temperatures, respectively (Boja & Girma, 2022). It is a vertisol soil. For the ginger, September to January is considered the monocropping season (main season). Although farmers also produce horticultural crops, pulses, and oil crops, the Ginir district is highly suited for the production of cereals.

According to the Central Statistical Agency's population projection, as cited in Boja *et al.*, (2022), the district's total population by 2021 was projected to be 203,751 (103,592 males and 100,159 females). The district's topography lies between 1200 and 2406 meters above sea level.

Data from the district agricultural office show that the district's land configuration is classified as plain, making up about 85% of the total, followed by mountainous areas (3%), rugged and gorge areas (12%), and hills and valleys (approximately 15%). In the same way, the district's land use data shows that 30.5% of the land is arable or cultivable, 31.2% is pasture, 35.6% is forest, and 2.7% is deemed to be swampy, mountainous, or otherwise unusable (GDAB, 2022).

## **Material used for compost preparation**

Different crop residues that were readily available in the area were used to make compost, including grasses, sorghum, haricot bean, wheat, teff, and maize straws, as well as a combination of straws and grass for bedding. An equal quantity of farmyard manure was added to each substrate. To prepare the pit for composting, the gathered substrates were chopped and added. Biomass from Parthenium weeds, crop residue, and farmyard manure from the experimental site were the materials used in this

experiment. Parthenium weed was gathered early in the rainy season, just before flowering and chopped into tiny pieces no larger than 2.5 cm. To keep the necessary C/N ratio in the process, wheat straw was employed as a good source of carbon. Similarly, other organic wastes, like ash, were employed to make better-balanced compost and utilize the waste. Parthenium to wheat straw ratio = 1:2.78 and Parthenium to cow dung ratio = 1:27.78 was the total amount and combination ratio of materials used in the formation of compost. To speed up the composting process, all of the green biomass from the parthenium weed was freshly harvested and chopped into small pieces.

### **Compost preparation**

A 1 m × 1 m × 1 m pit was prepared in the home garden of the farmer. To keep the stacking process at 60% moisture, water was sprayed. They were kept in a semi-aerobic environment and had a top layer plastered with a mixture of soil, dung, and wheat straw. A turning was performed after a month, and the moisture content was kept constant. When the pit was built in the shade, good quality compost was produced in 45 to 60 days at the ideal temperatures and rates of decomposition. The compost unit is built using materials that will last for the duration of the process. Before they reached flowering, the parthenium plants were gathered, mechanically chopped into 1–2-inch pieces, and allowed to decompose for around 20 days with agricultural and animal waste. The climate was ideal, and regular watering kept the temperature and moisture levels stable. Once the materials had been combined for 20 days, the temperature was recorded. Compost preparation was done using pit composting techniques, and the composting process lasted for sixty days.

### **Parthenium Compost Laboratory analysis**

Parthenium Samples of compost were taken from every compost pit. Following sifting, the samples were examined in soil laboratories at the Baatuu and Sinana Agricultural Research Centers to determine the compost quality. Using a pH meter and electrical conductivity, the pH and EC of compost were determined in the supernatant suspension of a 1:2.5 soil-to-water ratio (Rhoades, 1982). Utilizing Walkley and Black (1934) to calculate organic carbon. Bremner and Mulvaney's (1982) Kjeldahl method was used to calculate total nitrogen.

Total exchangeable bases (Ca<sup>2+</sup>, Mg<sup>2+</sup>, K<sup>+</sup>, and Na<sup>+</sup>) were measured using a flame photometer, atomic

absorption spectrometry (AAS) for Mg<sup>2+</sup> and Ca<sup>2+</sup>, and flame photometry for K<sup>+</sup> and Na<sup>+</sup> (Okalebo *et al.*, 2002). The method used to calculate cation exchange capacity (CEC) was (Chapman, 1965). Araya *et al.*, (2015) conducted a germination test on 100 seeds that were selected from the compost and planted in beds to determine which ones were viable.

## **Results and Discussion**

### **Certain chemical characteristics of parthenium compost**

#### **Electrical conductivity and pH**

The results of the laboratory analysis showed that the combination of farm yard manure and parthenium biomass with animal manure and wheat straw produced the highest (pH value) and lowest (pH value), respectively (Table 1). This result is consistent with that of Jouquet *et al.*, (2013), who reported that the pH range for compost was 6.8–8.41. According to research conducted by Spiers and Fietje (2000) and Araya *et al.*, (2015), a higher pH is associated with a higher K level, which is what causes high Electrical Conductivity (EC). The entire range of electrical conductivity (Ec) values, which range from 0.000056 to 0.000062 (dS/m) ds/m (Table 1), did not show any significant variation. According to (Santamaria *et al.*, 2001) and (Malagueta *et al.*, 2022) EC values of *Parthenium* compost were free from salinity. This slight increase in potassium ions (K<sup>+</sup>) and other ions during the course of decomposition may be the cause of the EC increase. According to Huang *et al.*, (2004), the breakdown of organic materials may release mineral salts like phosphates and ammonium ions, which could explain the rise in EC.

#### **Organic matter, Cation Exchange Capacity (CEC), and C: N Ratio**

According to the analyzed results, the compost made from Parthenium compost plus farm yard manure had the lowest mean values of organic carbon (35.2%). In comparison, the compost made from Parthenium compost plus crop residue had the highest mean value of organic matter (37.8%) (Table 1). When compared to the availability of organic matter in garden soil, the organic content of all types of compost is generally high. This result is consistent with research conducted by Mulugeta *et al.*, (2022); Tadele *et al.*, (2020). Each type of parthenium compost showed a low C:N ratio. A low C:N ratio suggests a higher mineralization rate. The results

show that parthenium compost made from parthenium biomass plus crop residue had the lowest percentage (11%) while parthenium compost made from farm yard manure plus crop residue had the highest percentage (11.4%) (Table 1). This result is in line with the findings of two other authors, Derib *et al.*, (2016) and Mulugeta *et al.*, (2022), who reported that vermicompost had a lower C:N ratio than regular compost. The parthenium compost produced using all treatment methods had a very high CEC, ranging from 34.8 to 53.2 cmol+ kg-1. This

outcome supported the research by Mulugeta *et al.*, (2022), which discovered that conventional compost contained 33.23 to 65.43 cmol+ kg-1 of CEC.

Higher concentrations of EC, OC, NT, and CEC were found in the compost made from parthenium combined with farm yard manure and crop residue. Veena and Shivani's (2012) study also demonstrated that parthenium, being a plant high in protein, is beneficial for soil and animal feed.

**Table.1** Organic matter and some macronutrients

Trt	pH-H <sub>2</sub> O (1:2.5)	EC (dS/m)	OM (%)	TN (%)	CEC (cmo(+)/kg)	C: N
<b>T1</b>	7.17	0.000062	35.2	1.83	34.8	11.2
<b>T2</b>	7.26	0.000056	36.1	1.91	46.2	11
<b>T3</b>	7.27	0.000058	37.8	1.98	53.2	11.4


**T1** = Parthenium compost + farm yard manure; **T2** = Parthenium compost + crop residue, **T3** = Parthenium compost + farm yard manure + crop residue

**Table.2** Exchangeable Basic Cations

Trt	Exchangeable Basic cations (cmol (+)/kg)				PBS (%)
	Ca	Mg	K	Na	
<b>T1</b>	4.56	1.30	1.51	0.24	21.87
<b>T2</b>	4.80	2.00	1.68	0.31	19.1
<b>T3</b>	5.40	3.25	1.86	0.33	20.38

Where **T1** = Parthenium compost + farm yard manure; **T2** = Parthenium compost + crop residue, **T3** = Parthenium compost + farm yard manure + crop residue

**Figure.1** Map of the study area.





## Total Nitrogen

In this study, the Parthenium weed's major nutrient composition was estimated to be lowest from Parthenium compost plus farm yard manure (1.83%), highest from Parthenium compost combined farm yard manure plus crop residue (1.98%), and nitrogen was recorded from the compost that was made. Findings from related research by Biradar *et al.*, (2005); Araya *et al.*, (2015) and Ameta *et al.*, (2016).

## Parthenium compost's exchangeable bases (Ca, Mg, K, and Na)

As was determined by Table 2, the results of the analysis indicated that the values for exchangeable bases (Ca, Mg, K, and Na) varied from 4.56 to 5.40 (col (+)/kg), 1.30 to 3.25 (col (+)/kg), 1.51 to 1.86 (col (+)/kg), and 0.24 to 0.33 (col (+)/kg), respectively. The Parthenium biomass combination of wheat straw and farm yard manure yielded the highest value in all cases, comparatively speaking than compost made from Parthenium biomass plus animal manure or Parthenium biomass plus wheat straw. In agreement with this Channappagoudar *et al.*, (2007) finding Compared to parthenium compost plus farm yard manure compost, the parthenium compost made with a combination of farm yard manure and other crop residue was generally richer in exchangeable cations. The outcome was in line with Amir and Fouzia's (2011) findings, which showed that parthenium compost made from mixed farmyard manure and parthenium biomass considerably increased the exchangeable bases (Ca, Ma, and K).

## Conclusions and Recommendations

These days, parthenium (*Parthenium hysterophorus*) is extensively dispersed throughout the agroecosystem and has emerged as a significant threat to land productivity and agricultural output. Although numerous attempts have been made by the government and various NGOs to stop or slow its growth, no appreciable shift has yet been seen. Composting parthenium weed is a novel way to extract maximum benefit and, as a result, limit the weed's spread. In terms of macro and micronutrients, compost is preferable to farm yard manure. They contribute significantly to the fertility of the soil and raise crop yields. Composting parthenium, can be used as an organic manure that effectively stops its alarming spread. The current study identified methods for controlling weeds and for using environmentally friendly technologies to support sustainable crop production and

soil productivity. The compost made from parthenium biomass, when combined with crop residue and farm yard manure, had higher nutrient contents than compost made solely from parthenium biomass. In general, it should be advised to raise public awareness, particularly among farmers, of the impact of *Parthenium hysterophorus* on agricultural productivity, ecosystem health, and management strategies. More research is generally required to determine the optimal rate of Parthenium compost application and how it affects crop yields and the physical and chemical characteristics of soil in field settings.

## Acknowledgement

We sincerely thank the members of the Sinana Agricultural Research Center's Soil Fertility Improvement and Soil and Water Conservation research team for their enthusiastic participation in this experiment. We also thank the Oromia Agricultural Research Institute for funding the project and the Sinana Agricultural Research Center for facilitating logistical services.

## Conflicts of Interest

No conflict of interest was disclosed by the author.

## References

- Ameta S K, Benjamin S, Ameta R, Ameta S C. Vishishta Composting: A Fastest Method and Eco-Friendly Recipe for Preparing Compost from *Parthenium hysterophorus* Weed. J Earth Environ Health Sci2016;2:103-8.
- Amir Khan and Fouzia Ishaq. 2011. Chemical nutrient analysis of different composts (Vermicompost and Pitcompost) and their effect on the growth of a vegetative crop *Pisum sativum*. Asian Journal of Plant Science and Research,1(1):116-130.
- Anbalagan M., Manivannan S. (2012): Assessment of the impact of invasive weed *Parthenium hysterophorus* L. mixed with organic supplements on growth and reproduction preference of *Lampito mauritii* (Kinberg) through vermitechnology. Int.J.Envirion.Biol.2(2):88-91.
- Ayele Shashie, Nigatu, Lisanework., Tana, Tamado., and Adkins, S. W. 2014. Impact of parthenium weed (*Parthenium hysterophorus* L.) on the above-ground and soil seed bank communities of range lands in Southeast Ethiopia. Global.

- Biradar, C. M., Saran, S., Raju, P. L. N. and Royb, P. S. (2005) Forest Canopy Density Stratification: How Relevant Is Biophysical Spectral Response Modelling Approach? Geocarto International, 20,1-7.
- Boja, M.; Girma, Z.; Dalle, G. Impacts of *Parthenium hysterophorus* L. on Plant Species Diversity in Ginir District, Southeastern Ethiopia. Diversity 2022, 14, 675.
- Bremner, J. M. and Mulvaney, C. S.(1982) Nitrogen-Total. In: Methods of soil analysis. Part 2. Chemical and microbiological properties, Page, A. L., Miller, R. H. and Keeney, D. R. Eds., American Society of Agronomy, Soil Science Society of America, Madison, Wisconsin, 595-624.
- Bruce, R. C., and Rayment, G. E. (1982). Analytical methods and interpretations used by the Agricultural Chemistry Branch for Soil and Land Use Surveys. Queensland Department of Primary Industries. BulletinQB8(2004), Indooroopilly, Queensland.
- Channappa Gowda B B, Biradar N R, Patil J B, Gasimani C A. Utilization of weed biomass as an organic source in sorghum. Karnataka JAgricSci2007;20:245-8.
- Charman, P. E. V., and Roper, M. M. (2007). Soil organic matter. In 'Soils—their properties and management'. 3rd edn. (Eds P.E.V. Charman and B. W. Murphy.) pp.276–285.(Oxford University Press:Melbourne.).
- DeribKifle, Gemechu Shumi and Abera Degefa. 2017. Characterization of Vermicompost for Major Plant Nutrient Contents and Manuring Value. Journal of Science and Sustainable Development,5(2),97-108.
- Devi Y N, Dutta B K, Sagolshemcha R, Singh N I. Allelopathic effect of *Parthenium hysterophorus* L. on growth and productivity of *Zea mays* L. and its phytochemical screening. IntJCurrMicrobiolApplSci2014;3:837-46.
- Evans H C (1997). *Parthenium hysterophorus*: A review of its weed status and the possibilities for biological control. Biocon.N.Info.18:89-98.
- Ginnir distinct agricultural office, 2022.
- Hailu Araya, Mitiku Haile, Arefayne Asmelash, Sue Edwards, and Tewolde Berhan Gebre Egziabher, 2015. Overcoming the Challenge of *Parthenium hysterophorus* through composting journal of agricultural science and food technology vol.1(6),pp.72-77.
- Hazelton, P. and Murphy, B. (2007) Interpreting Soil Test Results: What Do All the Numbers Mean? CSIRO Publishing, Collingwood. <https://doi.org/10.1071/9780643094680>.
- Jelin J., Dhanarajan M. S. (2013): Comparative physicochemical analysis of degrading *Parthenium (Parthenium hysterophorus)* and sawdust by a new approach to accelerating the composting rate. Int.J.Chem.Envirn.Biol.Sci.1(3):535-537.
- Kohli R K, Batish D R, Singh H P, Dogra K (2006). Status, invasiveness, and environmental threats of three tropical American invasive Weeds (*Parthenium hysterophorus* L., *Ageratum conyzoides* L., *Lantana camara* L.). BiologicalInvasions8:1501-1510.
- Kumari, P., Sahu, P. K., Soni, M. Y. and Awasthi, P. 2014. Determination of Organic Matter in Soil. SoilScience,37,549-556.
- Mesfin Boja, Nigusu Girma. The Perception of Local Peoples About *Parthenium hysterophorus* invasion and its impacts on plant biodiversity in ginir district, Southeastern Ethiopia. International Journal of Natural Resource Ecology and Management. Vol.7, No.1, 2022, pp.42-53. <https://doi.org/10.11648/j.ijnrem.20220701.16>.
- Metson, A. J. (1961). Methods of chemical analysis for soil survey samples. Soil Bureau Bulletin No.12, New Zealand Department of Scientific and Industrial Research, pp.168–175. (Government Printer: Wellington, New Zealand.).
- Mulugeta Eshetu, Daniel Abegeja, Tilahun Chibsa, Negash Bedaso. 2022 Worm Collection and Characterization of Vermicompost produced using different worm species and waste feeds materials at Sinanaon–Station of Bale highland southeastern Ethiopia. International Journal of Environmental and Agriculture Research;8(2).
- Njorge, J. M. 1991. Tolerance of *Bidens pilosa* and *Parthenium hysterophorus* L. Toparaquat (gramaxone) in Kenya coffee. Kenya Coffee:56:999-1001.
- Okalebo, J. R., Gathua, K. W. and Woome, P. L. (2002) Laboratory Methods for Soil and Plant Analysis: A Working Manual.TSBF, Nairobi.
- Rhoades, J. D. (1982) Cation Exchanges Capacity. In: Page, A. L., Buxton, R. H. and Miller Keeney, D. R., Eds., Methods of Soil Analysis, AmSocAgron, Madison,149-158.
- Santamaria R. S., Ferrera C. R, Almaraz S. J. J., Galvis S. A., Barois B. I. 2001. Dynamics and

- relationships among microorganisms, C-organic and N-total during composting and vermicomposting. *Agrociencia-Montecillo*,5(4),377-384.
- SeifuW/Kidan (1990) *Parthenium hysterophorus* L. A Recently Introduced Noxious Weed to Ethiopia. A Preliminary Reconnaissance Survey Report on Eastern Ethiopia. East Hararge Ministry of Agriculture, Ethiopia.
- Spiers T M, Fietje G (2000). Green waste compost as a component in soilless growing media. *CompostSci.Util*.8(1):19-23.
- Tadele Geremu, Habtamu Hailu, Alemayhu Diriba. 2020. Evaluation of Nutrient Content of Vermicompost Made from Different Substrates at Mechara Agricultural Research Center on Station, West Hararghe Zone, Oromia, Ethiopia. *Ecology and Evolutionary Biology*.5(4),125-130.
- Tamado T, Milberg P (2000). Weed florainarable fields of eastern Ethiopia with emphasis on the occurrence of *Parthenium hysterophorus*. *WeedRes*.40:507-521.
- Tefera T (2002). Allelopathic effects of *Parthenium hysterophorus* extracts on seed germination and seedling growth of *Eragrostis tef*. *J.Agron.CropSci*.188:306-310.
- Veena B K, Shivani M (2012). Biological utilities of *Parthenium hysterophorus*. *J.Appl.NaturalSci*.4(1):137-143.
- Wabuye, E., Lusweti, A., Bisikwa, J., Kyenune, G., Clark, K., Lotter, W. D., McConnachie, A. J. and Wondi, M., 2014. A roadside survey of the invasive weed *Parthenium hysterophorus* (Asteraceae) in East Africa. *Journal of East African Natural History*,103(1),49-57.
- Walkey, A. and Black, I. A Walkey, A. and Black, I. A. (1934) <https://doi.org/10.3390/d14080675>
- Yadav. 2015 Assessment of different organic supplements for degradation of *Parthenium hysterophorus* by vermitechnology *Journal of Environmental Health Science & Engineering*13:4.

**How to cite this article:**

Tesfaye Ketema Defar, Mulugeta Eshetu Diriba, Girma Getachew Gemechu and Regassa Gosa Telila. 2024. Determination of the Major Plant Nutrient Contents in *Parthenium (Parthenium hysterophorus* L.) Compost at Ginir District of Bale Zone, Oromia Region, Southeast Ethiopia. *Int.J.Curr.Res.Aca.Rev.* 12(4), 36-42. doi: <https://doi.org/10.20546/ijcrar.2024.1204.005>