



doi: <https://doi.org/10.20546/ijcrar.2021.906.001>

## Effect of Electroculture an *Arachis hypogaea* (L)

**R. U. Bhagyawant and M. B. Patil\***

*Department of Botany, Late Ramesh Warpudkar Arts, commerce and Science College, Sonpeth, Dist. Parbhani, India*

*\*Corresponding author*

### Abstract

Experiments were carried out to study effect of electriculture on Groundnut, during these experiments response of groundnut to 3V, 6V and 9V electricity was studied. It was found that various growth parameters like Height of plant, Number of elongation pegs, Number of leaves had shown significant response to the application of electricity. Yield Parameters like Number of pods per plant, Hundred seed weight, Seed size, Estimated weight was significantly increased due to application of electricity. Nutritional value of the groundnut seed was also studied plant applied with 3V and 9V electricity had shown decrease in fat percentage. There was no significant effect on fibre, Carbohydrate and protein percentage of groundnut seeds. Post-harvest soil estimation showed that majority of soil components were recorded in large quantity in experimental fields than the control field. Thus application of electricity was helpful in the retention of soil components.

### Article Info

Accepted: 25 May 2021

Available Online: 20 June 2021

### Keywords

Electroculture, Growth parameters, Yield Parameters Soil composition, *Arachishypogaea*.

### Introduction

Electroculture is a technique that uses electricity to improve seed germination, plant growth, quantity and yield. This technique is found useful to protect plants from disease and pests. Electromagnetic field influence physiological activity (Jaffe and Nuccitelli, 1977).

It is found that electroculture influence root growth (Souda *et al.*, 1970) the electric current also change the growth direction (Wanget *et al.*, 1989), (Filek *et al.*, 2003), enhance or inhibit flowering and stimulate embryogenesis (Dijak *et al.*, 1986) seed germination (Borisov, 1994; Yalgared M *et al.*, 2003; Panarese, 2009; Patil, 2018; Bhagyawant and Patil, 2021). Zhurbitskii and Shidlovskaya (1967) studied effect of electricity on uptake of irons in a solution that can affect absorption and incorporation of heavy metal iron such type of

results also reported by Murr 1964; Kotaka *et al.*, 1965; Krueger *et al.*, 1994.

Groundnut is one of the major oil-yielding crops, it is widely grown in the tropic and subtropic countries of the world. It is cultivated in 108 countries on about 22.2 million hectares. In India it is cultivated about 8 million hectares area. But due to high production cost and low productivity percentage area under cultivation of groundnut is decreasing. Present investigations are deals with the study of effect of electroculture on growth parameters, Yield parameters, Nutritional value, Soil component and oil percentage of the Groundnut.

### Materials and Methods

Experiment was carried out in the field located at Sonpeth Dist. Parbhani (19°1'34''N latitude and

76°28'18''E longitude at the elevation of 437m above the sea level). The experiment was conducted in complete randomized block designs. The field was prepared for sowing by following standard pre-sowing activities. The experimental field was divided into four equal plots of 10'x10' size. Three plots were supplied with 3V, 6V and 9V electricity, one plot was used as control. All the parameters except electricity was kept constant in all the fields.

Seeds were sown following standard method for groundnut cultivation. In experimental plots seeds were sown in between two electrodes. Each electrode was connected with the help of wire and finally it was connected to 3V/6V/9V DC(direct current)source of electricity.

Electricity treatment was given was given daily at morning (8am) for 10 min till harvesting.

Morphological and Yield parameters were studied ninety days after sowing. Height of the plant and length of root was measured in centimeters using scale and number of leaves, number of pegs, number of root nodules and number of pods were counted numerically. Seed size was measured by method describe by Wankhede *et al.*, (2019). Estimated yield was calculated quintal/acre by multiplying with 4.356 to the actual weight of pods yielded from the experimental plots.

Protein and fiber content of the seeds was estimated by the methods described by AOAC, for carbohydrate content estimated by method described by FAO while fat content was estimated by following IS 3579-1966 Reaffirmed 2016 method.

Soil samples were analyzed before sowing and after harvesting by methods described by FAO 2008.

The observations were recorded and tabulated in tables. Data was analyzed statistically using ANOVA.

## Results and Discussion

### Growth Parameters

Number of leaves per plant was increased significantly with the application of electricity. Maximum number of leaves (133.76) per plant was recorded in plants applied with 3V electricity; it is followed by plants applied with 6V electricity (110.64) and 9V (106.08).Control showed less number of leaves (66.72) similar results was found

by Gabdrakhmanova D and Qussiny, C (2011) in sweet pepper, Kanimarani (2018) in Cowpea cultivars and Afrasiyab *et al.*, (2020) in Chickpea.

In Control height of the plant was (28.77cm), application of electricity to the plants increased height of the plants. It was maximum in the plants applied with 3V electricity (34.18cm) followed by the plant applied with 9V electricity(31.12cm). The plants applied with 6V electricity showed an increase in the height of plant but it was statically non-significant. Increased height due to application of electricity may be due to increased cell division or cell elongation as a result of increased Gibberellins. Similar results were recorded by Dayal *et al.*, (1983) in tomato; Lakshmappa *et al.*, (2011) in wheat, bengal gram, green gram and moth bean; Laith Mohammad Shamma (2012) in faba bean.

Application of electricity increased root length in groundnut but these this increase in root length was statistically non-significant in plants applied with 3V and 6V electricity. Root length was maximum in plants applied with 9V electricity 54.02cm which was significant at P = 0.05.

In groundnut elongation pegs are produced after fertilization. The number of pegs per plant increased with the application of electricity the maximum number of pegs per plant were recorded in plants applied with 9V electricity (74.80) followed by the plants applied with 3V (67.7) and 6V (63.70) electricity. In control it was (56.40).

This increase in the number of pegs per plants due to application of electricity was statistically significant. This increase in the number of pegs indicates increased pollination due to application of the electricity.

Number of root nodules per plant was increased due to application of 3 V electricity (51.90) followed by the plant with the 9 Vs (48.70). It was decreased (43.50) due to application of 6 Vs electricity. But this change in the number of root nodules per plant was statistically non-significant. Thus application electricity has no significant role on root nodulation.

Seed size (l × b × t) in control was (241.39mm). The plants applied with 6V electricity showed maximum seed size (393.02mm) followed by plants applied with 9V (346.00mm) and 3 V (310.91mm) electricity. Electricity has played important role to increase seed size, it was statistically highly significant.

**Table.1** Effect of Electroculture on Growth Parameters of Groundnut plant

Character	Control	3 V	6V	9V	SE	CD 5%	CD1%
Number of leaves	66.72	133.76**	110.64**	106.08**	12.42	24.84	33.04
Height of plant (Cm)	28.77	34.18**	30.67 <sup>NS</sup>	31.12*	1.35	2.69	3.58
Length of root (Cm)	29.83	38.46 <sup>NS</sup>	38.92 <sup>NS</sup>	54.02*	10.35	20.71	27.54
No of peg	56.40	67.7**	63.70*	74.80**	3.36	6.88	9.29
No of root nodules	48.70	51.90 <sup>NS</sup>	43.50 <sup>NS</sup>	48.70 <sup>NS</sup>	3.41	7.00	9.45

**Table.2** Effect of Electroculture on Yield Parameters of Groundnut plant

Character	control	3 V	6V	9V	SE	CD 5%	CD1%
Number of pods/plant	40.44	64.64*	40.44 <sup>NS</sup>	52.84*	3.27	6.54	8.70
100 seed weight (gm)	43.57	50.32*	49.28 <sup>NS</sup>	55.24*	2.07	6.59	12.11
Seed size (length) mm	12.76	13.32 <sup>NS</sup>	14.12**	13.18 <sup>NS</sup>	0.31	0.63	0.84
Breadth(b) mm	8.24	8.86**	9.66**	9.29	0.24	0.48	0.64
Thickness(t) mm	7.19	7.97**	8.41**	8.46**	0.24	0.49	0.65
(l×b×t)	241.39	310.91**	393.02**	346.00**			
Estimated yield (Quintal/Acre)	22.99	41.94*	23.08 <sup>NS</sup>	32.67 <sup>NS</sup>	3.93	12.8	22.93

**Table.3** Effect of Electroculture on Nutritional Value of Groundnut seeds.

Nutrient	Control	3V	6V	9V	SE	CD5%	CD1%
Total fiber (%)	8.19	10.57 <sup>NS</sup>	7.72 <sup>NS</sup>	17.38*	1.93	6.13	11.27
Total carbohydrate (%)	16.01	11.35 <sup>NS</sup>	16.01 <sup>NS</sup>	14.32 <sup>NS</sup>	0.95	3.03	5.56
Total protein (%)	26.73	31.89 <sup>NS</sup>	27.09 <sup>NS</sup>	22.05 <sup>NS</sup>	1.74	5.54	10.17
Total fat (%)	46.9	43.96*	47.21 <sup>NS</sup>	43.73*	0.81	2.56	4.70

\*significant at p=0.05; \*\*significant at p=0.01; NS-non significant

**Table.4** Effect of Electroculture on Soil Components

Sr. No.	Nutrient	Pre-Sowing	Post-Harvest			
			CONTROL	3V	6V	9V
1	Nitrogen (Kg/ha)	188.16	150.33	213.25	163.07	125.44
2	Phosphorous (Kg/ha)	8.51	61.44	103.67	92.04	101.43
3	Potassium (Kg/ha)	873.6	599.20	862.40	752.64	976.64
4	Calcium (meq % )	30.77	31.49	34.56	26.88	30.72
5	Magnesium (meq %)	21.43	4.28	5.95	9.52	3.57
6	Sulphur (ppm)	67.5	292.00	277.00	281.00	209.00
7	Copper (ppm)	1.134	8.674	10.566	11.302	11.618
8	Iron (ppm)	0.260	1.209	0.190	0.205	0.205
9	Zinc (ppm)	0.346	0.426	1.084	0.926	0.998
10	Molybdenum (ppm)	0.094	0.097	0.105	0.077	0.075

Number of pods per plant increased with the application of electricity, maximum number of pods per plant(64.64) was recorded in plants applied with 3V electricity followed by the plants applied with 9V electricity (52.84) as compared to control(40.44) it was increased significantly. This might be due to stimulation of increased metabolism due to application of electricity.

Similar results were recorded by Lakshmappa (2011) in Bengal gram, green gram and moth bean, Laith Mohammed Shamma (2012) in faba bean.

Weight of 100 seeds was measured, in control it was (43.57gm). The application of electricity increased seed weight. Maximum weight of seeds(55.24gm) was found

in plants applied with 9V electricity followed by the plant applied with 3V electricity (50.32gm) and 6V electricity (49.28gm).

Increase in the 100 seeds weight was highly significant in plants applied with 3V and 9V electricity while increase in seed weight was in plants applied with 6V electricity was statistically non-significant.

Total yield (q/acre) was increased with the application of electricity; the plants applied with 3V electricity gave maximum yield (41.94q/acre) which was statistically significant. In plants applied with 9V electricity and 6V electricity yield was 32.67 q/acre and 32.08 q/acre respectively though it was increased it was statistically non significant.

As there was increase in the number of pods, 100 seeds weight and seed size significantly due to application of 3V electricity yield was also significant in the same field. Similar results were found in Francis *et al.*, (1989) in sunflower; Lale *et al.*, (2004) in cotton, Georgiy (2013) in wheat, barley and sugar beet, and Patil (2018) in *Raphanus*.

Nutritional analysis of groundnut seeds belonging to each field was carried out in order to study effect of electricity on nutritional value of groundnut. It was found that electricity has role to change carbohydrate and protein percentage of groundnut but it was statistically non-significant.

Application of 9V electricity increased fiber content total fat% significantly while application of 3V and 9V electricity decreased fat percentage in groundnut seeds significantly. As chemicals are consumed and released by the plants in its rhizosphere, cultivation of plants result in to the change in the soil nutrients.

Efforts were made to study effect of electricity on the soil nutrients in groundnut cultivated field. Before sowing 188.16 kg/ha soil nitrogen was present. Post-harvest soil estimation from the field applied with 3V electricity showed 213.25kg/ha nitrogen followed by field applied with 6V electricity 163.07kg/ha which was greater than the control(150.53kg/ha).

Thus application of electricity may have increased more nitrogen fixation in the soil. Same effect was seen on the majority of soil components. Thus application of electricity was helpful in the retention of soil components.

## References

- Afrasiyab A., Zafar J., Muhmmad H. (2020) Effect of electric field on seed germination and growth parameters of chickpea *Cicer arietinum* L. Ukrainian Journal of Ecology 10(4):12-16.
- Association of official analytical collaboration international (AOAC) official method 2011.11, chapter no.4.34,4.2.11.
- Bhagyawant R U and Patil, M B(2021). Effect of Electroculture on seed germination of certain crop plants. *Bioscience Discovery*, 12(2):69-72
- Borisov A M (1994). Activating the growing and development of plants by ionic-radiant processing; New materials and technology; thesis of research conf.Mosco.p.163.
- Dayal, S., Srivastava, K. G. and Singh, R. P. (1983). Growth Responses of tomato to seed and seedling exposure to Electrostatic field. *Indian J. Agric. Sci.* 53: 962-970.
- Dijak M, Sumith D L, Willson T J, Brown DCW.(1986).Stimulation of direct embryogenesis from mesophyll protoplast of *Medicagosativa*. *Plant Cell Report* 5:468-470.
- FAO Traditional methods and recent innovation, chapter no.2 method no.2.3
- Filek M, Biesage-Koscieliniak J, Barcinska I, Krekule J, Machackova I, Dubert F.(2003).The effect of electric current on flowering o grafted scion of non-vernalized winter rape. *Biological Plantarum* 46:625-628.
- Francis, M. S., Mathew, K. M., Subramanian, M. and Hariharan, M. (1989). Effect of pre-treatment of sunflower seeds With different intensities of electric current. *ActaBotanicaIndica* 17: 268-270.
- Gabdrakhmanova, D. and Qussiny C. 2011. Plantricity: The Effect of a Direct Electric Current on the Germination of Seeds and Growth of Seedlings. California State Science Fair.
- Georgiy, B. I. (2013). Impact of electromagnetic energy on the increasing yield capacity and growth stimulation of plants. *Ann. of Warsaw Uni. of Life Sci., Agric., no. 62 (Agric. and Forest Engg.):* 31–35
- Jaffe LF,Nucitelli R(1977). Electrical control of development. *Annual Review of Biophysics and Bioeneginnering* 6:445-476.
- Kotaka, S., Krueger, A. P., Nishizawa, K., Ohuchi, T., Takenobu, M., Kogure, Y., and Andriese, P. C. (1965a): Air ion effects on the Oxygen

- consumption of barley seedlings. Nature (Lond.), 208:1112-1113.
- Krueger A. P., S. Kotaka, and P. C. Andrise(1994). Effect of air containing  $O_2^-$ ,  $O_2$ ,  $CO_2^+$  on the growth of seedling of *Hordeum Vulguris*. Int.J.Biometerol.8.17.
- Laith M J AL-Shamma (2015) Variation of Faba Beans (*Vicia faba* L.) Traits Induced By Heat, Electric Shock and Mutagen Nitrous Acid. Baghdad Science Journal 12 (1) 80-88.
- Lakshmappa Ragha, Seema Mishra, V. Ramachandran, Manmohan Singh Bhatia (2011) Effects of Low-Power Microwave Fields on Seed Germination and Growth Rate. Journal of Electromagnetic Analysis and Application. (3)165-171.
- Lale, E., Mustafayev, S. A. and Fatih, K. (2004). Stimulative effect of high voltage electrical current on earliness, yield and fiber quality of cotton (*Gossypium hirsutum* L.). Pakistan J. of Biol. Sci. 7(4): 494&502.
- Murr, L. E.(1964). Mechanism of plant cell damage in electrostatic field. Nature 206,467-470.
- Official Methods of Analysis: AOAC International 20<sup>th</sup> edition;(2016)
- Panarese, V. (2009). Barley germination: Metabolic activity and effects of pulsed electric field. MSc Thesis. Lund University, Sweden.
- Patil, M B, (2018). Effect of Electroculture on seed germination and growth of *Raphanus sativus*(L). African Journal of Plant Science. 12(12), 350-353. <https://doi.org/10.5897/AJPS2018.1716>
- Sawsan Mohammed –saeed Ali Kanimarani (2018) Effect of Seed Exposure to Direct Electrical Current on Germination and Seedlings Growth of three Cowpeas (*Vigna unguiculata* L.) Cultivars. ZJPAS (2018),30 (5); 168-179
- Souda M, Toko K, Hayashi K, Ezaki S, Fujiyoshi T, Yamafuji K.(1990). Relationship between growth and electric oscillation in bean roots. Plant Physiology 93:532-536.
- Wang C, Rathmore K S, Robinson K R.(1989). The response of pollen to applied electrical fields. Developmental Biology136:405-410.
- Wankhade, V. R., R. C. Verma, B. L. Salvi, N. Jain and Jain, H. K. (2019). Assessment of Engineering Properties of Aonla Stone and Seeds. Int.J.Curr.Microbiol.App.Sci. 8(4): 2415-2423. Doi: <https://doi.org/10.20546/ijcmas.2019.804.281>
- Yalgared M, Mortzavi S A, Tabatabaie F(2008)Application of ultra sonic waves as a priming technique for accelerating and enhancing the germination of barley seed: optimization of method by the Taguchi approach.J.Inst.Brew.114(1):14-21.
- Zhurbitskii, Z. I., and Shidlovskaya, I. L. (1967): The action of an electrical field and ionized air on the absorption of mineral ions by wheat sprouts. Abstracted in Bio. Abstr., 1969, 100181.

**How to cite this article:**

Bhagyawant, R. U. and Patil, M. B. 2021. Effect of Electroculture an *Arachis hypogaea* (L). Int.J.Curr.Res.Aca.Rev. 9(06), 1-5. doi: <https://doi.org/10.20546/ijcrar.2021.906.001>